

No-Calc. Part

1. If  $y = \cos^2(2x)$ , then  $\frac{dy}{dx} =$

- (A)  $2 \cos 2x \sin 2x$  (C)  $2 \cos 2x$   
(B)  $-4 \sin 2x \cos 2x$  (D)  $-2 \cos 2x$   
(E)  $4 \cos 2x$

2. Of the following, the limit which exists is

- (A)  $\lim_{x \rightarrow 0} \left( \frac{1}{x^2} \right)$  (B)  $\lim_{x \rightarrow 0} \left( \frac{|x|}{x} \right)$  (C)  $\lim_{x \rightarrow 0} \left( \frac{1}{\sin x} \right)$   
(D)  $\lim_{x \rightarrow 0} (\ln x)$  (E)  $\lim_{x \rightarrow 0} \left( \frac{x^2 + 2x}{x} \right)$

3. The slope of the line tangent to the graph of  $y = \ln \sqrt{x}$  at  $(e^2, 1)$  is

- (A)  $\frac{e^2}{2}$  (B)  $\frac{2}{e^2}$  (C)  $\frac{1}{2e^2}$  (D)  $\frac{1}{2e}$  (E)  $\frac{1}{e}$

4. Which of the following functions is both continuous and differentiable at all  $x$  in the interval  $-2 \leq x \leq 2$ ?

- (A)  $f(x) = |x^2 - 1|$  (C)  $f(x) = \sqrt{x^2 + 1}$   
(B)  $f(x) = \sqrt{x^2 - 1}$  (D)  $f(x) = \frac{1}{x^2 - 1}$

5. Find the point on the graph of  $y = \sqrt{x}$  between  $(1, 1)$  and  $(9, 3)$  at which the tangent to the graph has the same slope as the line through  $(1, 1)$  and  $(9, 3)$ .

- (A)  $(1, 1)$  (C)  $(3, \sqrt{3})$   
(B)  $(2, \sqrt{2})$  (D)  $(4, 2)$   
(E) none of the above

6. Consider the function  $f(x) = \frac{x^4}{2} - \frac{x^5}{10}$ . The derivative of  $f$  attains its maximum value at  $x =$

- (A) 3 (C) 5  
(B) 4 (D) 0  
(E) there is no maximum

7. The acceleration,  $a(t)$ , of a body moving in a straight line is given in terms of time  $t$  by  $a(t) = 4 - 6t$ . If the velocity of the body is 20 at  $t = 0$  and if  $s(t)$  is the distance of the body from the origin at time  $t$ , what is  $s(3) - s(1)$ ?

- (A) -10 (C) 10 (E) 30  
(B) 0 (D) 20



14.  $\int_1^2 \frac{x^2 - x}{x^3} dx =$

- (A)  $\ln 2 - \frac{1}{2}$       (B)  $\ln 2 + \frac{1}{2}$       (C)  $\frac{1}{2}$       (D) 0      (E)  $\frac{1}{4}$

15. The edge of a cube is increasing at the uniform rate of 0.2 inches per second. At the instant when the total surface area becomes 150 square inches, what is the rate of increase, in cubic inches per second, of the volume of the cube?

- (A) 5 in<sup>3</sup>/sec  
 (B) 10 in<sup>3</sup>/sec  
 (C) 15 in<sup>3</sup>/sec  
 (D) 20 in<sup>3</sup>/sec  
 (E) 25 in<sup>3</sup>/sec

16.  $\int_0^{\sqrt{3}} \frac{x dx}{\sqrt{1+x^2}} =$

- (A)  $\frac{1}{2}$       (B) 1      (C) 2      (D)  $\ln 2$   
 (E)  $\text{Arctan } 2 - \frac{\pi}{4}$

17. Which of the following is true about the graph of  $f(x) = \ln|x^2 - 4|$  in the interval  $(-2, 2)$ ?

- (A)  $f$  is increasing.  
 (B)  $f$  attains a relative minimum at  $(0, 0)$ .  
 (C)  $f$  has a range of all real numbers.  
 (D)  $f$  is concave down.  
 (E)  $f$  has an asymptote at  $x = 0$ .

18. If  $g(x) = \text{Arcsin } 2x$ , then  $g'(x) =$

- (A)  $2\text{Arccos } 2x$       (B)  $2 \csc 2x \cot 2x$       (C)  $\frac{2}{1+4x^2}$   
 (D)  $\frac{2}{\sqrt{4x^2-1}}$       (E)  $\frac{2}{\sqrt{1-4x^2}}$

19.  $\int x(x^2 - 1)^4 dx =$

- (A)  $\frac{1}{10}(x^2)(x^2 - 1)^5 + C$       (C)  $\frac{1}{5}(x^3 - x)^5 + C$   
 (B)  $\frac{1}{10}(x^2 - 1)^5 + C$       (D)  $\frac{1}{5}(x^2 - 1)^5 + C$   
 (E)  $\frac{1}{5}(x^2 - x)^5 + C$

20. If  $y = e^{kx}$ , then  $\frac{d^5 y}{dx^5} =$

- (A)  $k^5 e^x$       (C)  $5! e^{kx}$   
 (B)  $k^5 e^{kx}$       (D)  $5! e^x$   
 (E)  $5e^{kx}$


21. The graph of  $f$  is shown at the right. Which of the following statements are true?

I.  $f(2) > f'(1)$

II.  $\int_0^1 f(x) dx > f'(3.5)$

III.  $\int_{-1}^1 f(x) dx > \int_{-1}^2 f(x) dx$

the graph of  $f$



- (A) I only    (B) II only    (C) I and II only    (D) II and III only    (E) I, II, III

22. If  $g(x) = \sqrt{x}(x-1)^{2/3}$ , then the domain of  $g'$  is

(A)  $\{x \mid 0 < x\}$

(C)  $\{x \mid 0 < x < 1 \text{ or } x > 1\}$

(B)  $\{x \mid x \neq 0 \text{ and } x \neq 1\}$

(D)  $\{x \mid 0 < x < 1\}$

(E)  $\{x \mid \text{all real numbers}\}$

23. A point moves on the  $x$ -axis so that its distance from the origin at time  $t$  is given by  $10t - 4t^2$ . What is the *total* distance covered by the point between  $t = 1$  and  $t = 2$ ?

(A) 1.0


(C) 2.0

(B) 1.5

(D) 2.5

(E) 3.0

24. At which point on the graph of  $y = g(x)$  below is  $g'(x) = 0$  and  $g''(x) = 0$ ?



(A) A

(B) B

(C) C

(D) D

(E) E

25. If  $y$  is a differentiable function of  $x$ , then the slope of the tangent to the curve  $xy - 2y + 4y^2 = 6$  at the point where  $y = 1$  is

(A)  $\frac{1}{12}$

(B)  $-\frac{1}{10}$

(C)  $-\frac{1}{6}$

(D)  $\frac{1}{4}$

(E)  $-\frac{5}{6}$

26. The area of the region bounded above by  $y = 1 + \sec^2 x$ , below by  $y = 0$ , on the left by  $x = 0$  and on the right by  $x = \frac{\pi}{4}$  is approximately

(A) 1 units<sup>2</sup>

(B) 1.25 units<sup>2</sup>

(C) 1.5 units<sup>2</sup>

(D) 1.75 units<sup>2</sup>

(E) 2 units<sup>2</sup>

27. A solution of the equation  $\frac{dy}{dx} + 2xy = 0$  that contains the point  $(0, e)$  is

(A)  $y = e^{1-x^2}$

(C)  $y = e^{1-x}$

(E)  $y = e^{x^2}$

(F)  $y = e^{1+x^2}$

(D)  $y = e^{1+x}$

I.  $F(1) = 0$       II.  $F'(1) = 0$        $F''(1) = 1$

(A) I and II only

(C) II and III only

(B) I and III only

(D) I, II, III

(E) none

$$F(x) = \int_1^x \ln(2t-1) dt$$

### Calculator Part:

1. How many points of inflection does the graph of  $y = \cos x + \frac{1}{3} \cos 3x - \frac{1}{5} \cos 5x$  have on the interval  $0 \leq x \leq \pi$ ?

- (A) 1
- (B) 2
- (C) 3
- (D) 4
- (E) 5

2. Oil is leaking from a tanker at the rate of  $R(t) = 500e^{-0.2t}$  gallons per hour, where  $t$  is measured in hours. The amount of oil that has leaked out after 10 hours is closest to

- (A) 2140 gals
- (B) 2150 gals
- (C) 2160 gals
- (D) 2170 gals
- (E) 2180 gals

3. The sale of lumber  $S$  (in millions of square feet) for the years 1980 to 1990 is modeled by the function


$$S(t) = 0.46 \cos(0.45t + 3.15) + 3.4$$

where  $t$  is the time in years with  $t = 0$  corresponding to the beginning of 1980. Determine the year when lumber sales were increasing at the greatest rate.

- (A) 1982
- (B) 1983
- (C) 1984
- (D) 1985
- (E) 1986

4. The graph of  $f$  over the interval  $[1, 9]$  is shown in the figure. Using the data in the figure, find a midpoint approximation with 4 equal subdivisions for

$$\int_1^9 f(x) dx.$$



- (A) 20
- (B) 21
- (C) 22
- (D) 23
- (E) 24

5. Let the base of a solid be the first quadrant region enclosed by the  $x$ -axis and one arch of the graph of  $y = \sin x$ . If all cross sections perpendicular to the  $x$ -axis are squares, then the volume of the solid is approximately

- (A)  $0.52 \text{ units}^3$       (C)  $1.05 \text{ units}^3$   
 (B)  $0.79 \text{ units}^3$       (D)  $1.57 \text{ units}^3$   
 (E)  $2.00 \text{ units}^3$

6. If  $f(x) = 2x + \sin x$  and the function  $g$  is the inverse of  $f$ , then  $g'(2) =$

- (A) 0.32  
 (B) 0.34  
 (C) 0.36  
 (D) 0.38  
 (E) 0.40

7. Administrators at Massachusetts General Hospital believe that the hospital's expenditures  $E(B)$ , measured in dollars, are a function of how many beds  $B$  are in use with

$$E(B) = 14000 + (B + 1)^2.$$

On the other hand, the number of beds  $B$  is a function of time  $t$ , measured in days, and it is estimated that

$$B(t) = 20 \sin\left(\frac{t}{10}\right) + 50.$$


At what rate are the expenditures decreasing when  $t = 100$ ?

- (A) 120 dollars/day      (C) 130 dollars/day  
 (B) 125 dollars/day      (D) 135 dollars/day  
 (E) 140 dollars/day

8. Let  $f$  be a function that has domain  $[-2, 5]$ . The graph of  $f'$  is shown at the right. Which of the following statements are TRUE?

- I.  $f$  has a relative maximum at  $x = -1$ .  
 II.  $f$  has an absolute minimum at  $x = 0$ .  
 III.  $f$  is concave down for  $-2 < x < 0$ .  
 IV.  $f$  has inflection points at  $x = 0$  and  $x = 2$  and  $x = 3$ .

the graph of  $f'$



- (A) I, II, IV      (B) I, III, IV      (C) II, III, IV      (D) I, II, III      (E) I, II, III, IV

9. On which interval is the graph of  $f(x) = 4x^{3/2} - 3x^2$  both concave down and increasing?

- (A)  $(0, 1)$   
 (B)  $\left(0, \frac{1}{2}\right)$       (D)  $\left(\frac{1}{4}, \frac{1}{2}\right)$   
 (C)  $\left(0, \frac{1}{4}\right)$       (E)  $\left(\frac{1}{4}, 1\right)$

Graph of  $y = f(x)$  is defined for all  $x \geq 0$ , contains the point  $(0, 1)$ , has  $\frac{dy}{dx} = 3\sqrt{xy}$  and  $f(x) > 0$  for all  $x$ , then  $f(x) =$

- (A)  $(x^3 + 1)^2$  (C)  $(x^{1/2} + 1)^2$   
(B)  $(x^{3/2} + 1)^2$  (D)  $(x^3 - 1)^2$   
(E)  $(x^{3/2} - 1)^2$

11. If  $\sin 3x - 1 = \int_a^x f(t) dt$ , then the value of  $a$  is

- (A) 0 (C) -1  
(B) 1 (D)  $\frac{\pi}{3}$  (E)  $\frac{\pi}{6}$

12. If  $xy^2 = 20$  and  $x$  is decreasing at the rate of 3 units per second, the rate at which  $y$  is changing when  $y = 2$  is nearest to

- (A) -0.6 units/sec  
(B) -0.2 units/sec  
(C) 0.2 units/sec  
(D) 0.6 units/sec  
(E) 1.0 units/sec

13. An approximation for  $\int_{-1}^2 e^{\sin(1.5x-1)} dx$  using a right-hand Riemann sum with three equal subdivisions is nearest to

- (A) 2.5 (C) 4.5 (E) 6.5  
(B) 3.5 (D) 5.5

14. If  $f(x)$  is defined on  $-\pi \leq x \leq \pi$  and  $\frac{dy}{dx} = \frac{\cos x}{x^2 + 1}$ , which of the following statements about the graph of  $y = f(x)$  is true?


- (A) The graph has no relative extremum.  
(B) The graph has one point of inflection and two relative extrema.  
(C) The graph has two points of inflection and one relative extremum.  
(D) The graph has two points of inflection and two relative extrema.  
(E) The graph has three points of inflection and two relative extrema.

15. The graph of the function  $f$  is shown at the right. If the function  $G$  is defined by

$$G(x) = \int_{-4}^x f(t) dt, \text{ for } -4 \leq x \leq 4, \text{ which of}$$

the following statements about  $G$  are true?

- I.  $G$  is increasing on  $(1, 2)$ .  
 II.  $G$  is decreasing on  $(-4, -3)$ .  
 III.  $G(0) < 0$ .



- (A) None      (B) II only      (C) III only      (D) II and III only      (E) I and II only

16. The function  $f$  is defined on all the reals such that  $f(x) = \begin{cases} x^2 + kx - 3 & \text{for } x \leq 1 \\ 3x + b & \text{for } x > 1. \end{cases}$

For which of the following values of  $k$  and  $b$  will the function  $f$  be both continuous and differentiable on its entire domain?

- (A)  $k = -1, b = -3$   
 (B)  $k = 1, b = 3$   
 (C)  $k = 1, b = 4$   
 (D)  $k = 1, b = -4$   
 (E)  $k = -1, b = 6$

17. A particle moves along the  $x$ -axis with velocity at time  $t$  given by:  $v(t) = t + 2 \sin t$ . If the particle is at the origin when  $t = 0$ , its position at the time when  $v = 6$  is  $x =$

- (A) 17.14      (B) 19.16      (C) 23.18      (D) 29.20      (E) 39.30