

P A R T I I

C H A P T E R 6

Applications of Integration

Section 6.1	Area of a Region Between Two Curves	264
Section 6.2	Volume: The Disk Method	271
Section 6.3	Volume: The Shell Method	278
Section 6.4	Arc Length and Surfaces of Revolution	282
Section 6.5	Work	287
Section 6.6	Moments, Centers of Mass, and Centroids	290
Section 6.7	Fluid Pressure and Fluid Force	297
Review Exercises		299
Problem Solving		305

C H A P T E R 6

Applications of Integration

Section 6.1 Area of a Region Between Two Curves

Solutions to Even-Numbered Exercises

$$2. A = \int_{-2}^2 [(2x + 5) - (x^2 + 2x + 1)] dx = \int_{-2}^2 (-x^2 + 4) dx$$




$$4. A = \int_0^1 (x^2 - x^3) dx$$

$$6. A = 2 \int_0^1 [(x - 1)^3 - (x - 1)] dx$$

$$8. \int_{-1}^1 [(1 - x^2) - (x^2 - 1)] dx$$

$$10. \int_2^3 \left[\left(\frac{x^3}{3} - x \right) - \frac{x}{3} \right] dx$$

$$12. \int_{-\pi/4}^{\pi/4} (\sec^2 x - \cos x) dx$$




$$14. f(x) = 2 - \frac{1}{2}x$$

$$g(x) = 2 - \sqrt{x}$$

$$A \approx 1$$

Matches (a)




$$16. A = \int_2^8 \left[\left(10 - \frac{1}{2}x \right) - \left(-\frac{3}{8}x(x - 8) \right) \right] dx$$

$$= \int_2^8 \left(\frac{3}{8}x^2 - \frac{7}{2}x + 10 \right) dx$$


$$= \left[\frac{x^3}{8} - \frac{7x^2}{4} + 10x \right]_2^8$$

$$= (64 - 112 + 80) - (1 - 7 + 20) = 18$$




18. The points of intersection are given by

$$\begin{aligned} -x^2 + 4x + 1 &= x + 1 \\ -x^2 + 3x &= 0 \\ x^2 = 3x \text{ when } x = 0, 3 \\ A &= \int_0^3 [(-x^2 + 4x + 1) - (x + 1)] dx \\ &= \int_0^3 (-x^2 + 3x) dx \\ &= \left[-\frac{x^3}{3} + \frac{3x^2}{2} \right]_0^3 \\ &= -9 + \frac{27}{2} = \frac{9}{2} \end{aligned}$$



22. $A = \int_1^5 \left(\frac{1}{x^2} - 0 \right) dx = \left[-\frac{1}{x} \right]_1^5 = \frac{4}{5}$





24. The points of intersection are given by

$$\begin{aligned} \sqrt[3]{x-1} &= x-1 \\ x-1 &= (x-1)^3 = x^3 - 3x^2 + 3x - 1 \\ x^3 - 3x^2 + 2x &= 0 \\ x(x^2 - 3x + 2) &= 0 \\ x(x-2)(x-1) &= 0 \Rightarrow x = 0, 1, 2 \\ A &= 2 \int_0^1 [(x-1) - \sqrt[3]{x-1}] dx \\ &= 2 \left[\frac{x^2}{2} - x - \frac{3}{4}(x-1)^{4/3} \right]_0^1 \\ &= 2 \left[\left(\frac{1}{2} - 1 - 0 \right) - \left(-\frac{3}{4} \right) \right] = \frac{1}{2} \end{aligned}$$

20. The points of intersection are given by:

$$\begin{aligned} -x^2 + 4x + 2 &= x + 2 \\ x(3-x) &= 0 \text{ when } x = 0, 3 \end{aligned}$$

$$\begin{aligned} A &= \int_0^3 [f(x) - g(x)] dx \\ &= \int_0^3 [(-x^2 + 4x + 2) - (x + 2)] dx \\ &= \int_0^3 (-x^2 + 3x) dx = \left[-\frac{x^3}{3} + \frac{3}{2}x^2 \right]_0^3 = \frac{9}{2} \end{aligned}$$




26. The points of intersection are given by:


$$2y - y^2 = -y$$

$$y(y - 3) = 0 \text{ when } y = 0, 3$$


$$\begin{aligned} A &= \int_0^3 [f(y) - g(y)] dy \\ &= \int_0^3 [(2y - y^2) - (-y)] dy \\ &= \int_0^3 (3y - y^2) dy = \left[\frac{3}{2}y^2 - \frac{1}{3}y^3 \right]_0^3 = \frac{9}{2} \end{aligned}$$



$$\begin{aligned} 28. A &= \int_0^3 [f(y) - g(y)] dy \\ &= \int_0^3 \left[\frac{y}{\sqrt{16 - y^2}} - 0 \right] dy \\ &= -\frac{1}{2} \int_0^3 (16 - y^2)^{-1/2} (-2y) dy \\ &= \left[-\sqrt{16 - y^2} \right]_0^3 = 4 - \sqrt{7} \approx 1.354 \end{aligned}$$



$$\begin{aligned} 30. A &= \int_0^1 \left(4 - \frac{4}{2-x} \right) dx \\ &= \left[4x + 4 \ln |2-x| \right]_0^1 \\ &= 4 - 4 \ln 2 \\ &\approx 1.227 \end{aligned}$$




32. The point of intersection is given by:

$$x^3 - 2x + 1 = -2x$$

$$x^3 + 1 = 0 \text{ when } x = -1$$

$$\begin{aligned} A &= \int_{-1}^1 [f(x) - g(x)] dx \\ &= \int_{-1}^1 [(x^3 - 2x + 1) - (-2x)] dx \\ &= \int_{-1}^1 (x^3 + 1) dx = \left[\frac{x^4}{4} + x \right]_{-1}^1 = 2 \end{aligned}$$

Numerical Approximation: 2.0




34. The points of intersection are given by:

$$x^4 - 2x^2 = 2x^2$$

$$x^2(x^2 - 4) = 0 \text{ when } x = 0, \pm 2$$

$$\begin{aligned} A &= 2 \int_0^2 [2x^2 - (x^4 - 2x^2)] dx \\ &= 2 \int_0^2 (4x^2 - x^4) dx \\ &= 2 \left[\frac{4x^3}{3} - \frac{x^5}{5} \right]_0^2 = \frac{128}{15} \end{aligned}$$

Numerical Approximation: 8.533




36. $f(x) = x^4 - 4x^2$, $g(x) = x^3 - 4x$

The points of intersection are given by:

$$x^4 - 4x^2 = x^3 - 4x$$

$$x^4 - x^3 - 4x^2 + 4x = 0$$


$$x(x-1)(x+2)(x-2) = 0 \text{ when } x = -2, 0, 1, 2$$



$$\begin{aligned} A &= \int_{-2}^0 [(x^3 - 4x) - (x^4 - 4x^2)] dx + \int_0^1 [(x^4 - 4x^2) - (x^3 - 4x)] dx + \int_1^2 [(x^3 - 4x) - (x^4 - 4x^2)] dx \\ &= \frac{248}{30} + \frac{37}{60} + \frac{53}{60} = \frac{293}{30} \end{aligned}$$


Numerical Approximation: $8.267 + 0.617 + 0.883 \approx 9.767$

38. $A = \int_0^3 \left[\frac{6x}{x^2 + 1} - 0 \right] dx$
 $= \left[3 \ln(x^2 + 1) \right]_0^3$
 $= 3 \ln 10$
 ≈ 6.908




Numerical Approximation: 6.908


42. $A = \int_{-\pi/2}^{\pi/6} (\cos 2x - \sin x) dx$
 $= \left[\frac{1}{2} \sin 2x + \cos x \right]_{-\pi/2}^{\pi/6}$
 $= \left(\frac{\sqrt{3}}{4} + \frac{\sqrt{3}}{2} \right) - (0) = \frac{3\sqrt{3}}{4} \approx 1.299$



40. $A = \int_0^4 x \sqrt{\frac{4-x}{4+x}} dx \approx 3.434$




44. $A = \int_0^1 \left[(\sqrt{2} - 4)x + 4 - \sec \frac{\pi x}{4} \tan \frac{\pi x}{4} \right] dx$
 $= \left[\frac{\sqrt{2}-4}{2}x^2 + 4x - \frac{4}{\pi} \sec \frac{\pi x}{4} \right]_0^1$
 $= \left(\frac{\sqrt{2}-4}{2} + 4 - \frac{4}{\pi} \sqrt{2} \right) - \left(-\frac{4}{\pi} \right)$
 $= \frac{\sqrt{2}}{2} + 2 + \frac{4}{\pi}(1 - \sqrt{2}) \approx 2.1797$




46. From the graph we see that f and g intersect twice at $x = 0$ and $x = 1$.

$$\begin{aligned} A &= \int_0^1 [g(x) - f(x)] dx \\ &= \int_0^1 [(2x+1) - 3^x] dx \\ &= \left[x^2 + x - \frac{1}{\ln 3} (3^x) \right]_0^1 \\ &= 2 \left(1 - \frac{1}{\ln 3} \right) \approx 0.180 \end{aligned}$$




48. $A = \int_0^\pi [(2 \sin x + \cos 2x) - 0] dx$

$$= \left[-2 \cos x + \frac{1}{2} \sin 2x \right]_0^\pi = 4$$



$$\begin{aligned} \mathbf{50.} \quad A &= \int_1^5 \left[\frac{4 \ln x}{x} - 0 \right] dx \\ &= \left[2(\ln x)^2 \right]_1^5 = 2(\ln 5)^2 \approx 5.181 \end{aligned}$$




$$\mathbf{52. (a)} \quad y = \sqrt{x} e^x, \quad y = 0, \quad x = 0, \quad x = 1$$

$$\text{(b)} \quad A = \int_0^1 \sqrt{x} e^x dx.$$


No, it cannot be evaluated by hand.

$$\text{(c)} \quad 1.2556$$




$$\mathbf{54.} \quad F(x) = \int_0^x \left(\frac{1}{2}t^2 + 2 \right) dt = \left[\frac{1}{6}t^3 + 2t \right]_0^x = \frac{x^3}{6} + 2x$$


$$\text{(a)} \quad F(0) = 0$$



$$\text{(b)} \quad F(4) = \frac{4^3}{6} + 2(4) = \frac{56}{3}$$




$$\text{(c)} \quad F(6) = 36 + 12 = 48$$




$$\mathbf{56.} \quad F(y) = \int_{-1}^y 4e^{x/2} dx = \left[8e^{x/2} \right]_{-1}^y = 8e^{y/2} - 8e^{-1/2}$$


$$\text{(a)} \quad F(-1) = 0$$




$$\text{(b)} \quad F(0) = 8 - 8e^{-1/2} \approx 3.1478$$



$$\text{(c)} \quad F(4) = 8e^2 - 8e^{-1/2} \approx 54.2602$$



$$\begin{aligned} \mathbf{58.} \quad A &= \int_2^4 \left[\left(\frac{9}{2}x - 12 \right) - (x - 5) \right] dx + \int_4^6 \left[\left(-\frac{5}{2}x + 16 \right) - (x - 5) \right] dx \\ &= \int_2^4 \left(\frac{7}{2}x - 7 \right) dx + \int_4^6 \left(-\frac{7}{2}x + 21 \right) dx \\ &= \left[\frac{7}{4}x^2 - 7x \right]_2^4 + \left[-\frac{7}{4}x^2 + 21x \right]_4^6 = 7 + 7 = 14 \end{aligned}$$



60. $f(x) = \frac{1}{x^2 + 1}$

$$f'(x) = -\frac{2x}{(x^2 + 1)^2}$$

$$\text{At } \left(1, \frac{1}{2}\right), f'(1) = -\frac{1}{2}.$$

Tangent line:

$$y - \frac{1}{2} = -\frac{1}{2}(x - 1) \text{ or } y = -\frac{1}{2}x + 1$$

The tangent line intersects $f(x) = \frac{1}{x^2 + 1}$ at $x = 0$.

$$A = \int_0^1 \left[\frac{1}{x^2 + 1} - \left(-\frac{1}{2}x + 1 \right) \right] dx = \left[\arctan x + \frac{x^2}{4} - x \right]_0^1 = \frac{\pi - 3}{4} \approx 0.0354$$

62. Answers will vary. See page 417.

64. $x^3 \geq x$ on $[-1, 0]$


$$x^3 \leq x \text{ on } [0, 1]$$

Both functions symmetric to origin

$$\int_{-1}^0 (x^3 - x) dx = -\int_0^1 (x^3 - x) dx.$$

$$\text{Thus, } \int_{-1}^1 (x^3 - x) dx = 0.$$

$$A = 2 \int_0^1 (x - x^3) dx = 2 \left[\frac{x^2}{2} - \frac{x^4}{4} \right]_0^1 = \frac{1}{2}$$



66. Proposal 2 is better, since the cumulative deficit (the area under the curve) is less.

68. $A = 2 \int_0^9 (9 - x) dx = 2 \left[9x - \frac{x^2}{2} \right]_0^9 = 81$

$$2 \int_0^{9-b} [(9 - x) - b] dx = \frac{81}{2}$$



$$2 \int_0^{9-b} [(9 - b) - x] dx = \frac{81}{2}$$

$$2 \left[(9 - b)x - \frac{x^2}{2} \right]_0^{9-b} = \frac{81}{2}$$

$$(9 - b)(9 - b) = \frac{81}{2}$$

$$9 - b = \frac{9}{\sqrt{2}}$$


$$b = 9 - \frac{9}{\sqrt{2}} \approx 2.636$$



70. $\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n (4 - x_i^2) \Delta x$

where $x_i = -2 + \frac{4i}{n}$ and $\Delta x = \frac{4}{n}$ is the same as

$$\int_{-2}^2 (4 - x^2) dx = \left[4x - \frac{x^3}{3} \right]_{-2}^2 = \frac{32}{3}.$$



72.
$$\begin{aligned} \int_0^5 [(7.21 + 0.26t + 0.02t^2) - (7.21 + 0.1t + 0.01t^2)] dt &= \int_0^5 (0.01t^2 + 0.16t) dt \\ &= \left[\frac{0.01t^3}{3} + \frac{0.16t^2}{2} \right]_0^5 \\ &= \frac{29}{12} \text{ billion} \approx \$2,417 \text{ billion} \end{aligned}$$

74. 5% : $P_1 = 893,000 e^{(0.05)t}$

$\frac{1}{2}\%$: $P_2 = 893,000 e^{(0.035)t}$

Difference in profits over 5 years:

$$\begin{aligned} \int_0^5 [893,000e^{0.05t} - 893,000e^{0.035t}] dt &= 893,000 \left[\frac{e^{0.05t}}{0.05} - \frac{e^{0.035t}}{0.035} \right]_0^5 \\ &\approx 893,000[(25.6805 - 34.0356) - (20 - 28.5714)] \\ &\approx 893,000(0.2163) \approx \$193,156 \end{aligned}$$

Note: Using a graphing utility you obtain \$193,183.

76. The curves intersect at the point where the slope of y_2 equals that of y_1 , 1.

$$y_2 = 0.08x^2 + k \Rightarrow y'_2 = 0.16x = 1 \Rightarrow x = \frac{1}{0.16} = 6.25$$

(a) The value of k is given by

$$y_1 = y_2$$

$$6.25 = (0.08)(6.25)^2 + k$$

$$k = 3.125.$$

$$\begin{aligned} \text{(b) Area} &= 2 \int_0^{6.25} (y_2 - y_1) dx \\ &= 2 \int_0^{6.25} (0.08x^2 + 3.125 - x) dx \\ &= 2 \left[\frac{0.08x^3}{3} + 3.125x - \frac{x^2}{2} \right]_0^{6.25} \\ &= 2(6.510417) \approx 13.02083 \end{aligned}$$

78. (a) $A \approx 6.031 - 2 \left[\pi \left(\frac{1}{16} \right)^2 \right] - 2 \left[\pi \left(\frac{1}{8} \right)^2 \right] \approx 5.908$

(b) $V = 2A \approx 2(5.908) \approx 11.816 \text{ m}^3$

(c) $5000V \approx 5000(11.816) = 59,082 \text{ pounds}$

80. True

Section 6.2 Volume: The Disk Method

2. $V = \pi \int_0^2 (4 - x^2)^2 dx = \pi \int_0^2 (x^4 - 8x^2 + 16) dx = \pi \left[\frac{x^5}{5} - \frac{8x^3}{3} + 16x \right]_0^2 = \frac{256\pi}{15}$

4. $V = \pi \int_0^3 (\sqrt{9 - x^2})^2 dx = \pi \int_0^3 (9 - x^2) dx$
 $= \pi \left[9x - \frac{x^3}{3} \right]_0^3 = 18\pi$

6. $2 = 4 - \frac{x^2}{4}$
 $x^2 = 8$
 $x = \pm 2\sqrt{2}$
 $V = \pi \int_{-2\sqrt{2}}^{2\sqrt{2}} \left[\left(4 - \frac{x^2}{4}\right)^2 - (2)^2 \right] dx$
 $= 2\pi \int_0^{2\sqrt{2}} \left[\frac{x^4}{16} - 2x^2 + 12 \right] dx$
 $= 2\pi \left[\frac{x^5}{80} - \frac{2x^3}{3} + 12x \right]_0^{2\sqrt{2}}$
 $= 2\pi \left[\frac{128\sqrt{2}}{80} - \frac{32\sqrt{2}}{3} + 24\sqrt{2} \right]$
 $= \frac{448\sqrt{2}}{15}\pi \approx 132.69$


8. $y = \sqrt{16 - x^2} \Rightarrow x = \sqrt{16 - y^2}$
 $V = \pi \int_0^4 (\sqrt{16 - y^2})^2 dy = \pi \int_0^4 (16 - y^2) dy$
 $= \pi \left[16y - \frac{y^3}{3} \right]_0^4 = \frac{128\pi}{3}$

10. $V = \pi \int_1^4 (-y^2 + 4y)^2 dy = \pi \int_1^4 (y^4 - 8y^3 + 16y^2) dy$
 $= \pi \left[\frac{y^5}{5} - 2y^4 + \frac{16y^3}{3} \right]_1^4$
 $= \frac{459\pi}{15} = \frac{153\pi}{5}$

12. $y = 2x^2, y = 0, x = 2$


(a) $R(y) = 2, r(y) = \sqrt{y}/2$

$$V = \pi \int_0^8 \left(4 - \frac{y}{2}\right) dy = \pi \left[4y - \frac{y^2}{4} \right]_0^8 = 16\pi$$




(c) $R(x) = 8, r(x) = 8 - 2x^2$

$$\begin{aligned} V &= \pi \int_0^2 [64 - (64 - 32x^2 + 4x^4)] dx \\ &= \pi \int_0^2 (32x^2 - 4x^4) dx = 4\pi \int_0^2 (8x^2 - x^4) dx \\ &= 4\pi \left[\frac{8}{3}x^3 - \frac{1}{5}x^5 \right]_0^2 = \frac{896\pi}{15} \end{aligned}$$




(b) $R(x) = 2x^2, r(x) = 0$

$$V = \pi \int_0^2 4x^4 dx = \pi \left[\frac{4x^5}{5} \right]_0^2 = \frac{128\pi}{5}$$



(d) $R(y) = 2 - \sqrt{y/2}, r(y) = 0$


$$\begin{aligned} V &= \pi \int_0^8 \left(2 - \sqrt{\frac{y}{2}}\right)^2 dy \\ &= \pi \int_0^8 \left(4 - 4\sqrt{\frac{y}{2}} + \frac{y}{2}\right) dy \\ &= \pi \left[4y - \frac{4\sqrt{2}}{3}y^{3/2} + \frac{y^2}{4} \right]_0^8 = \frac{16\pi}{3} \end{aligned}$$



14. $y = 6 - 2x - x^2$, $y = x + 6$ intersect at $(-3, 3)$ and $(0, 6)$.


(a) $R(x) = 6 - 2x - x^2$, $r(x) = x + 6$

$$\begin{aligned} V &= \pi \int_{-3}^0 [(6 - 2x - x^2)^2 - (x + 6)^2] dx \\ &= \pi \int_{-3}^0 (x^4 + 4x^3 - 9x^2 - 36x) dx \\ &= \pi \left[\frac{1}{5}x^5 + x^4 - 3x^3 - 18x^2 \right]_{-3}^0 = \frac{243\pi}{5} \end{aligned}$$




(b) $R(x) = (6 - 2x - x^2) - 3$, $r(x) = (x + 6) - 3$

$$\begin{aligned} V &= \pi \int_{-3}^0 [(3 - 2x - x^2)^2 - (x + 3)^2] dx \\ &= \pi \int_{-3}^0 (x^4 + 4x^3 - 3x^2 - 18x) dx \\ &= \pi \left[\frac{1}{5}x^5 + x^4 - 3x^3 - 9x^2 \right]_{-3}^0 = \frac{108\pi}{5} \end{aligned}$$




16. $R(x) = 4 - \frac{x^3}{2}$, $r(x) = 0$

$$\begin{aligned} V &= \pi \int_0^2 \left(4 - \frac{x^3}{2} \right)^2 dx \\ &= \pi \int_0^2 \left[16 - 4x^3 + \frac{x^6}{4} \right] dx \\ &= \pi \left[16x - x^4 + \frac{x^7}{28} \right]_0^2 \\ &= \pi \left[32 - 16 + \frac{128}{28} \right] = \frac{144}{7}\pi \end{aligned}$$




18. $R(x) = 4$, $r(x) = 4 - \sec x$

$$\begin{aligned} V &= \pi \int_0^{\pi/3} [(4)^2 - (4 - \sec x)^2] dx \\ &= \pi \int_0^{\pi/3} (8 \sec x - \sec^2 x) dx \\ &= \pi \left[8 \ln|\sec x + \tan x| - \tan x \right]_0^{\pi/3} \\ &= \pi [(8 \ln|2 + \sqrt{3}| - \sqrt{3}) - (8 \ln|1 + 0| - 0)] \\ &= \pi [8 \ln(2 + \sqrt{3}) - \sqrt{3}] \approx 27.66 \end{aligned}$$




20. $R(y) = 6$, $r(y) = 6 - (6 - y) = y$

$$\begin{aligned} V &= \pi \int_0^4 [(6)^2 - (y)^2] dy \\ &= \pi \left[36y - \frac{y^3}{3} \right]_0^4 = \frac{368\pi}{3} \end{aligned}$$




22. $R(y) = 6 - \frac{6}{y}$, $r(y) = 0$

$$\begin{aligned} V &= \pi \int_2^6 \left(6 - \frac{6}{y}\right)^2 dy \\ &= 36\pi \int_2^6 \left(1 - \frac{2}{y} + \frac{1}{y^2}\right) dy \\ &= 36\pi \left[y - 2\ln|y| - \frac{1}{y}\right]_2^6 \\ &= 36\pi \left[\left(\frac{35}{6} - 2\ln 6\right) - \left(\frac{3}{2} - 2\ln 2\right)\right] \\ &= 36\pi \left(\frac{13}{3} + 2\ln \frac{1}{3}\right) = 12\pi(13 - 6\ln 3) \approx 241.59 \end{aligned}$$



26. $R(x) = \frac{3}{x+1}$, $r(x) = 0$


$$\begin{aligned} V &= \pi \int_0^8 \left(\frac{3}{x+1}\right)^2 dx \\ &= 9\pi \int_0^8 (x+1)^{-2} dx \\ &= 9\pi \left[-\frac{1}{x+1}\right]_0^8 = 8\pi \end{aligned}$$



$$\begin{aligned} 30. V &= \pi \int_0^4 \left[\left(4 - \frac{1}{2}x\right)^2 - (\sqrt{x})^2 \right] dx + \pi \int_4^8 \left[(\sqrt{x})^2 - \left(4 - \frac{1}{2}x\right)^2 \right] dx \\ &= \pi \int_0^4 \left(\frac{x^2}{4} - 5x + 16 \right) dx + \pi \int_4^8 \left(-\frac{x^2}{4} + 5x - 16 \right) dx \\ &= \pi \left[\frac{x^3}{12} - \frac{5x^2}{2} + 16x \right]_0^4 + \pi \left[-\frac{x^3}{12} + \frac{5x^2}{2} - 16x \right]_4^8 \\ &= \frac{88}{3}\pi + \frac{56}{3}\pi = 48\pi \end{aligned}$$



24. $R(x) = x\sqrt{4-x^2}$, $r(x) = 0$

$$\begin{aligned} V &= 2\pi \int_0^2 [x\sqrt{4-x^2}]^2 dx \\ &= 2\pi \int_0^2 (4x^2 - x^4) dx \\ &= 2\pi \left[\frac{4x^3}{3} - \frac{x^5}{5} \right]_0^2 \\ &= \frac{128\pi}{15} \end{aligned}$$



28. $R(x) = e^{x/2}$, $r(x) = 0$

$$\begin{aligned} V &= \pi \int_0^4 (e^{x/2})^2 dx \\ &= \pi \int_0^4 e^x dx \\ &= \left[\pi e^x \right]_0^4 \\ &= \pi(e^4 - 1) \approx 168.38 \end{aligned}$$



32. $y = 9 - x^2$, $y = 0$, $x = 2$, $x = 3$


$$x = \sqrt{9 - y}$$

$$V = \pi \int_0^5 [(\sqrt{9 - y})^2 - 2^2] dy$$

$$= \pi \int_0^5 (5 - y) dy$$


$$= \pi \left[5y - \frac{y^2}{2} \right]_0^5$$

$$= \pi \left(25 - \frac{25}{2} \right) = \frac{25\pi}{2}$$



34. $V = \pi \int_0^{\pi/2} [\cos x]^2 dx \approx 2.4674$


36. $V = \pi \int_1^3 [\ln x]^2 dx \approx 3.2332$



38. $V = \pi \int_0^5 [2 \arctan(0.2x)]^2 dx \approx 15.4115$


40. $A \approx \frac{3}{4}$

Matches (b)




42. $V = \int_a^b A(x) dx \quad \text{or} \quad V = \int_c^d A(y) dy$


44. (a)



(b)



(c)




$$a < c < b.$$

46. $R(x) = \frac{r}{h}x$, $r(x) = 0$

$$V = \pi \int_0^h \frac{r^2}{h^2} x^2 dx$$


$$= \left[\frac{r^2 \pi}{3h^2} x^3 \right]_0^h$$

$$= \frac{r^2 \pi}{3h^2} h^3 = \frac{1}{3} \pi r^2 h$$



48. $x = \sqrt{r^2 - y^2}$, $R(y) = \sqrt{r^2 - y^2}$, $r(y) = 0$

$$\begin{aligned} V &= \pi \int_h^r (\sqrt{r^2 - y^2})^2 dy \\ &= \pi \int_h^r (r^2 - y^2) dy \\ &= \pi \left[r^2 y - \frac{y^3}{3} \right]_h^r \\ &= \pi \left[\left(r^3 - \frac{r^3}{3} \right) - \left(r^2 h - \frac{h^3}{3} \right) \right] \\ &= \pi \left(\frac{2r^3}{3} - r^2 h + \frac{h^3}{3} \right) \\ &= \frac{\pi}{3} (2r^3 - 3r^2 h + h^3) \end{aligned}$$



52. $y = \begin{cases} \sqrt{0.1x^3 - 2.2x^2 + 10.9x + 22.2}, & 0 \leq x \leq 11.5 \\ 2.95, & 11.5 < x \leq 15 \end{cases}$

$$\begin{aligned} V &= \pi \int_0^{11.5} (\sqrt{0.1x^3 - 2.2x^2 + 10.9x + 22.2})^2 dx + \pi \int_{11.5}^{15} 2.95^2 dx \\ &= \pi \left[\frac{0.1x^4}{4} - \frac{2.2x^3}{3} + \frac{10.9x^2}{2} + 22.2x \right]_0^{11.5} + \pi [2.95^2 x]_{11.5}^{15} \\ &\approx 1031.9016 \text{ cubic centimeters} \end{aligned}$$

54. (a) First find where $y = b$ intersects the parabola:

$$b = 4 - \frac{x^2}{4}$$

$$x^2 = 16 - 4b = 4(4 - b)$$

$$x = 2\sqrt{4 - b}$$

$$\begin{aligned} V &= \int_0^{2\sqrt{4-b}} \pi \left[4 - \frac{x^2}{4} - b \right]^2 dx + \int_{2\sqrt{4-b}}^4 \pi \left[b - 4 + \frac{x^2}{4} \right]^2 dx \\ &= \int_0^4 \pi \left[4 - \frac{x^2}{4} - b \right]^2 dx \\ &= \pi \int_0^4 \left[\frac{x^4}{16} - 2x^2 + \frac{bx^2}{2} + b^2 - 8b + 16 \right] dx \\ &= \pi \left[\frac{x^5}{80} - \frac{2x^3}{3} + \frac{bx^3}{6} + b^2 x - 8bx + 16x \right]_0^4 \\ &= \pi \left[\frac{64}{5} - \frac{128}{3} + \frac{32}{3}b + 4b^2 - 32b + 64 \right] = \pi \left[4b^2 - \frac{64}{3}b + \frac{512}{15} \right] \end{aligned}$$

50. (a) $V = \pi \int_0^4 (\sqrt{x})^2 dx = \pi \int_0^4 x dx = \left[\frac{\pi x^2}{2} \right]_0^4 = 8\pi$

Let $0 < c < 4$ and set

$$\pi \int_0^c x dx = \left[\frac{\pi x^2}{2} \right]_0^c = \frac{\pi c^2}{2} = 4\pi.$$

$$c^2 = 8$$

$$c = \sqrt{8} = 2\sqrt{2}$$

Thus, when $x = 2\sqrt{2}$, the solid is divided into two parts of equal volume.



(b) Set $\pi \int_0^c x dx = \frac{8\pi}{3}$ (one third of the volume). Then

$$\frac{\pi c^2}{2} = \frac{8\pi}{3}, \quad c^2 = \frac{16}{3}, \quad c = \frac{4}{\sqrt{3}} = \frac{4\sqrt{3}}{3}.$$

To find the other value, set $\pi \int_0^d x dx = \frac{16\pi}{3}$ (two thirds of the volume). Then


$$\frac{\pi d^2}{2} = \frac{16\pi}{3}, \quad d^2 = \frac{32}{3}, \quad d = \frac{\sqrt{32}}{\sqrt{3}} = \frac{4\sqrt{6}}{3}.$$

The x -values that divide the solid into three parts of equal volume are $x = (4\sqrt{3})/3$ and $x = (4\sqrt{6})/3$.



54. —CONTINUED—

(b) graph of $V(b) = \pi \left[4b^2 - \frac{64}{3}b + \frac{512}{15} \right]$



Minimum Volume is 17.87 for $b = 2.67$

(c) $V'(b) = \pi \left[8b - \frac{64}{3} \right] = 0 \Rightarrow b = \frac{64/3}{8} = \frac{8}{3} = 2\frac{2}{3}$

$V''(b) = 8\pi > 0 \Rightarrow b = \frac{8}{3}$ is a relative minimum.


56. (a) $V = \int_0^{10} \pi[f(x)]^2 dx$

Simpson's Rule: $b - a = 10 - 0 = 10$, $n = 10$

$$V \approx \frac{\pi}{3} [(2.1)^2 + 4(1.9)^2 + 2(2.1)^2 + 4(2.35)^2 + 2(2.6)^2 + 4(2.85)^2 + 2(2.9)^2 + 4(2.7)^2 + 2(2.45)^2 + 4(2.2)^2 + (2.3)^2]$$

$$\approx \frac{\pi}{3} [178.405] \approx 186.83 \text{ cm}^3$$


(b) $f(x) = 0.00249x^4 - 0.0529x^3 + 0.3314x^2 - 0.4999x + 2.112$



(c) $V \approx \int_0^{10} \pi f(x)^2 dx \approx 186.35 \text{ cm}^3$

58. $V = \frac{1}{2}(10)(2)(3) = 30 \text{ m}^3$


60.



Base of Cross Section = $2\sqrt{4 - x^2}$

(a) $A(x) = b^2 = (2\sqrt{4 - x^2})^2$

$$V = \int_{-2}^2 4(4 - x^2) dx \\ = 4 \left[4x - \frac{x^3}{3} \right]_{-2}^2 = \frac{128}{3}$$




(b) $A(x) = \frac{1}{2}bh = \frac{1}{2}(2\sqrt{4 - x^2})(\sqrt{3}\sqrt{4 - x^2})$

$$= \sqrt{3}(4 - x^2)$$

$$V = \sqrt{3} \int_{-2}^2 (4 - x^2) dx$$

$$= \sqrt{3} \left[4x - \frac{x^3}{3} \right]_{-2}^2 = \frac{32\sqrt{3}}{3}$$




—CONTINUED—

60. —CONTINUED—

$$(c) A(x) = \frac{1}{2}\pi r^2$$

$$= \frac{\pi}{2}(\sqrt{4-x^2})^2 = \frac{\pi}{2}(4-x^2)$$


$$V = \frac{\pi}{2} \int_{-2}^2 (4-x^2) dx = \frac{\pi}{2} \left[4x - \frac{x^3}{3} \right]_{-2}^2 = \frac{16\pi}{3}$$



$$(d) A(x) = \frac{1}{2}bh$$

$$= \frac{1}{2}(2\sqrt{4-x^2})(\sqrt{4-x^2}) = 4-x^2$$

$$V = \int_{-2}^2 (4-x^2) dx = \left[4x - \frac{x^3}{3} \right]_{-2}^2 = \frac{32}{3}$$




62. The cross sections are squares. By symmetry, we can set up an integral for an eighth of the volume and multiply by 8.

$$A(y) = b^2 = (\sqrt{r^2-y^2})^2$$

$$V = 8 \int_0^r (r^2 - y^2) dy$$

$$= 8 \left[r^2y - \frac{1}{3}y^3 \right]_0^r$$

$$= \frac{16}{3}r^3$$




$$64. V = \pi \int_{-\sqrt{R^2-r^2}}^{\sqrt{R^2-r^2}} [(\sqrt{R^2-x^2})^2 - r^2] dx$$

$$= 2\pi \int_0^{\sqrt{R^2-r^2}} (R^2 - r^2 - x^2) dx$$

$$= 2\pi \left[(R^2 - r^2)x - \frac{x^3}{3} \right]_0^{\sqrt{R^2-r^2}}$$

$$= 2\pi \left[(R^2 - r^2)^{3/2} - \frac{(R^2 - r^2)^{3/2}}{3} \right]$$

$$= \frac{4}{3}\pi(R^2 - r^2)^{3/2}$$




66. (a) When $a = 1$: $|x| + |y| = 1$ represents a square.

When $a = 2$: $|x|^2 + |y|^2 = 1$ represents a circle.

$$(b) |y| = (1 - |x|^a)^{1/a}$$

$$A = 2 \int_{-1}^1 (1 - |x|^a)^{1/a} dx = 4 \int_0^1 (1 - x^a)^{1/a} dx$$

To approximate the volume of the solid, form n slices, each of whose area is approximated by the integral above. Then sum the volumes of these n slices.



Section 6.3 Volume: The Shell Method

2. $p(x) = x$

$$h(x) = 1 - x$$

$$V = 2\pi \int_0^1 x(1-x) dx$$

$$= 2\pi \int_0^1 (x - x^2) dx = 2\pi \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = \frac{\pi}{3}$$

$$V = 2\pi \int_0^2 x(4-x^2) dx$$

4. $p(x) = x$

$$h(x) = 8 - (x^2 + 4) = 4 - x^2$$

$$V = 2\pi \int_0^2 (4x - x^3) dx$$


$$= 2\pi \left[2x^2 - \frac{x^4}{4} \right]_0^2 = 8\pi$$

6. $p(x) = x$

$$h(x) = \frac{1}{2}x^2$$

$$V = 2\pi \int_0^6 \frac{1}{2}x^3 dx$$

$$= \left[\frac{\pi x^4}{4} \right]_0^6 = 324\pi$$




8. $p(x) = x$

$$h(x) = 4 - x^2$$

$$V = 2\pi \int_0^2 (4x - x^3) dx$$

$$= 2\pi \left[2x^2 - \frac{1}{4}x^4 \right]_0^2 = 8\pi$$




10. $p(x) = x$

$$h(x) = 4 - 2x$$

$$V = 2\pi \int_0^2 x(4 - 2x) dx$$

$$= 2\pi \int_0^2 (4x - 2x^2) dx$$

$$= 2\pi \left[2x^2 - \frac{2}{3}x^3 \right]_0^2 = \frac{16\pi}{3}$$




12. $p(x) = x$

$$h(x) = \frac{\sin x}{x}$$

$$V = 2\pi \int_0^\pi x \left[\frac{\sin x}{x} \right] dx$$

$$= 2\pi \int_0^\pi \sin x dx = \left[-2\pi \cos x \right]_0^\pi = 4\pi$$



14. $p(y) = -y$ ($p(y) \geq 0$ on $[-2, 0]$)

$$h(y) = 4 - (2 - y) = 2 + y$$

$$V = 2\pi \int_{-2}^0 (-y)(2+y) dy$$

$$= 2\pi \int_{-2}^0 (-2y - y^2) dy$$

$$= 2\pi \left[-y^2 - \frac{y^3}{3} \right]_{-2}^0 = \frac{8\pi}{3}$$

16. $p(y) = y$


$$h(y) = 16 - y^2$$

$$V = 2\pi \int_0^4 y(16 - y^2) dy$$

$$= 2\pi \int_0^4 (16y - y^3) dy$$

$$= 2\pi \left[8y^2 - \frac{y^4}{4} \right]_0^4$$

$$= 2\pi[128 - 64] = 128\pi$$




18. $p(x) = 2 - x$

$$h(x) = 4x - x^2 - x^2 = 4x - 2x^2$$

$$V = 2\pi \int_0^2 (2-x)(4x-2x^2) dx$$

$$= 2\pi \int_0^2 (8x - 8x^2 + 2x^3) dx$$

$$= 2\pi \left[4x^2 - \frac{8}{3}x^3 + \frac{1}{2}x^4 \right]_0^2 = \frac{16\pi}{3}$$




20. $p(x) = 6 - x$

$$h(x) = \sqrt{x}$$

$$V = 2\pi \int_0^4 (6-x)\sqrt{x} dx$$

$$= 2\pi \int_0^4 (6x^{1/2} - x^{3/2}) dx$$

$$= 2\pi \left[4x^{3/2} - \frac{2}{5}x^{5/2} \right]_0^4 = \frac{192\pi}{5}$$



22. (a) Disk


$$R(x) = \frac{10}{x^2}, r(x) = 0$$

$$V = \pi \int_1^5 \left(\frac{10}{x^2} \right)^2 dx$$

$$= 100\pi \int_1^5 x^{-4} dx$$

$$= 100\pi \left[\frac{x^{-3}}{-3} \right]_1^5$$

$$= -\frac{100\pi}{3} \left[\frac{1}{125} - 1 \right] = \frac{496}{15}\pi$$



(b) Shell

$$R(x) = x, r(x) = 0$$

$$V = 2\pi \int_1^5 x \left(\frac{10}{x^2} \right) dx$$

$$= 20\pi \int_1^5 \frac{1}{x} dx$$

$$= 20\pi \left[\ln|x| \right]_1^5 = 20\pi \ln 5$$

(c) Disk

$$R(x) = 10, r(x) = 10 - \frac{10}{x^2}$$

$$V = \pi \int_1^5 \left[10^2 - \left(10 - \frac{10}{x^2} \right)^2 \right] dx$$

$$= \pi \left[\frac{100}{3x^3} - \frac{200}{x} \right]_1^5 = \frac{1904}{15}\pi$$

24. (a) Disk

$$R(x) = (a^{2/3} - x^{2/3})^{3/2}$$


$$r(x) = 0$$

$$V = \pi \int_{-a}^a (a^{2/3} - x^{2/3})^3 dx$$


$$= 2\pi \int_0^a (a^2 - 3a^{4/3}x^{2/3} + 3a^{2/3}x^{4/3} - x^2) dx$$


$$= 2\pi \left[a^2x - \frac{9}{5}a^{4/3}x^{5/3} + \frac{9}{7}a^{2/3}x^{7/3} - \frac{1}{3}x^3 \right]_0^a$$

$$= 2\pi \left(a^3 - \frac{9}{5}a^3 + \frac{9}{7}a^3 - \frac{1}{3}a^3 \right) = \frac{32\pi a^3}{105}$$



(b) Same as part a by symmetry






28. $2\pi \int_0^4 x \left(\frac{x}{2}\right) dx$


represents the volume of the solid generated by revolving the region bounded by $y = x/2$, $y = 0$, and $x = 4$ about the y -axis by using the Shell Method.

$$\pi \int_0^2 [16 - (2y)^2] dy = \pi \int_0^2 [(4)^2 - (2y)^2] dy$$


represents this same volume by using the Disk Method.



Disk Method



(b) $V = 2\pi \int_1^3 \frac{2x}{1 + e^{1/x}} dx \approx 19.0162$




(b) $V = 2\pi \int_0^1 x \sqrt{1 - x^3} dx \approx 2.3222$

34. $y = \tan x$, $y = 0$, $x = 0$, $x = \frac{\pi}{4}$

Volume ≈ 1

Matches (e)




36. Total volume of the hemisphere is $\frac{1}{2} \left(\frac{4}{3}\right) \pi r^3 = \frac{2}{3} \pi (3)^3 = 18\pi$. By the Shell Method, $p(x) = x$, $h(x) = \sqrt{9 - x^2}$. Find x_0 such that

$$\begin{aligned} 6\pi &= 2\pi \int_0^{x_0} x \sqrt{9 - x^2} dx \\ 6 &= -\int_0^{x_0} (9 - x^2)^{1/2} (-2x) dx \\ &= \left[-\frac{2}{3} (9 - x^2)^{3/2} \right]_0^{x_0} = 18 - \frac{2}{3} (9 - x_0^2)^{3/2} \end{aligned}$$

$$(9 - x_0^2)^{3/2} = 18$$

$$x_0 = \sqrt{9 - 18^{2/3}} \approx 1.460.$$

Diameter: $2\sqrt{9 - 18^{2/3}} \approx 2.920$



$$\begin{aligned}
38. \quad V &= 4\pi \int_{-r}^r (R-x)\sqrt{r^2-x^2} dx \\
&= 4\pi R \int_{-r}^r \sqrt{r^2-x^2} dx - 4\pi \int_{-r}^r x\sqrt{r^2-x^2} dx \\
&= 4\pi R \left(\frac{\pi r^2}{2} \right) + \left[2\pi \left(\frac{2}{3} \right) (r^2-x^2)^{3/2} \right]_{-r}^r \\
&= 2\pi^2 r^2 R
\end{aligned}$$

$$\begin{aligned}
40. \quad (a) \quad &\text{Area region} = \int_0^b [ab^n - ax^n] dx \\
&= \left[ab^n x - a \frac{x^{n+1}}{n+1} \right]_0^b \\
&= ab^{n+1} - a \frac{b^{n+1}}{n+1} \\
&= ab^{n+1} \left(1 - \frac{1}{n+1} \right) = ab^{n+1} \left(\frac{n}{n+1} \right) \\
R_1(n) &= \frac{ab^{n+1} \left(\frac{n}{n+1} \right)}{(ab^n)b} = \frac{n}{n+1} \\
(b) \quad &\lim_{n \rightarrow \infty} R_1(n) = \lim_{n \rightarrow \infty} \frac{n}{n+1} = 1 \\
&\lim_{n \rightarrow \infty} (ab^n)b = \infty
\end{aligned}$$

$$\begin{aligned}
(c) \quad &\textbf{Disk Method:} \\
V &= 2\pi \int_0^b x(ab^n - ax^n) dx \\
&= 2\pi a \int_0^b (xb^n - x^{n+1}) dx \\
&= 2\pi a \left[\frac{b^n}{2}x^2 - \frac{x^{n+2}}{n+2} \right]_0^b \\
&= 2\pi a \left[\frac{b^{n+2}}{2} - \frac{b^{n+2}}{n+2} \right] = \pi ab^{n+2} \left(\frac{n}{n+2} \right) \\
R_2(n) &= \frac{\pi ab^{n+2} \left(\frac{n}{n+2} \right)}{(\pi b^2)(ab^n)} = \left(\frac{n}{n+2} \right) \\
(d) \quad &\lim_{n \rightarrow \infty} R_2(n) = \lim_{n \rightarrow \infty} \left(\frac{n}{n+2} \right) = 1 \\
&\lim_{n \rightarrow \infty} (\pi b^2)(ab^n) = \infty
\end{aligned}$$

(e) As $n \rightarrow \infty$, the graph approaches the line $x = 1$.

$$\begin{aligned}
42. \quad (a) \quad &V = 2\pi \int_0^4 xf(x) dx \\
&= \frac{2\pi(40)}{3(4)} [0 + 4(10)(45) + 2(20)(40) + 4(30)(20) + 0] \\
&= \frac{20\pi}{3}[5800] \approx 121,475 \text{ cubic feet}
\end{aligned}$$

$$(b) \quad \text{Top line: } y - 50 = \frac{40 - 50}{20 - 0}(x - 0) = -\frac{1}{2}x \Rightarrow y = -\frac{1}{2}x + 50$$

$$\text{Bottom line: } y - 40 = \frac{0 - 40}{40 - 20}(x - 20) = -2(x - 20) \Rightarrow y = -2x + 80$$

$$\begin{aligned}
V &= 2\pi \int_0^{20} x \left(-\frac{1}{2}x + 50 \right) dx + 2\pi \int_{20}^{40} x(-2x + 80) dx \\
&= 2\pi \int_0^{20} \left(-\frac{1}{2}x^2 + 50x \right) dx + 2\pi \int_{20}^{40} (-2x^2 + 80x) dx \\
&= 2\pi \left[-\frac{x^3}{6} + 25x^2 \right]_0^{20} + 2\pi \left[-\frac{2x^3}{3} + 40x^2 \right]_{20}^{40} \\
&= 2\pi \left[\frac{26,000}{3} \right] + 2\pi \left[\frac{32,000}{3} \right] \\
&\approx 121,475 \text{ cubic feet}
\end{aligned}$$

(Note that Simpson's Rule is exact for this problem.)

Section 6.4 Arc Length and Surfaces of Revolution

2. (1, 2), (7, 10)

(a) $d = \sqrt{(7 - 1)^2 + (10 - 2)^2} = 10$

(b) $y = \frac{4}{3}x + \frac{2}{3}$

$$y' = \frac{4}{3}$$

$$s = \int_1^7 \sqrt{1 + \left(\frac{4}{3}\right)^2} dx = \left[\frac{5}{3}x\right]_1^7 = 10$$

6. $y = \frac{3}{2}x^{2/3} + 4$

$$y' = x^{-1/3}, [1, 27]$$

$$\begin{aligned} s &= \int_1^{27} \sqrt{1 + \left(\frac{1}{x^{1/3}}\right)^2} dx \\ &= \int_1^{27} \sqrt{\frac{x^{2/3} + 1}{x^{2/3}}} dx \\ &= \frac{3}{2} \int_1^{27} \sqrt{x^{2/3} + 1} \left(\frac{2}{3x^{1/3}}\right) dx \\ &= \left[\frac{3}{2} \cdot \frac{2}{3} (x^{2/3} + 1)^{3/2} \right]_1^{27} \\ &= 10^{3/2} - 2^{3/2} \approx 28.794 \end{aligned}$$

10. $y = \frac{1}{2}(e^x + e^{-x})$

$$y' = \frac{1}{2}(e^x - e^{-x}), [0, 2]$$


$$1 + (y')^2 = \left[\frac{1}{2}(e^x + e^{-x}) \right]^2, [0, 2]$$

$$s = \int_0^2 \sqrt{\left[\frac{1}{2}(e^x + e^{-x}) \right]^2} dx$$

$$= \frac{1}{2} \int_0^2 (e^x + e^{-x}) dx$$

$$= \frac{1}{2} \left[e^x - e^{-x} \right]_0^2 = \frac{1}{2} \left(e^2 - \frac{1}{e^2} \right) \approx 3.627$$

14. (a) $y = \frac{1}{1+x}, 0 \leq x \leq 1$



(b) $y' = -\frac{1}{(1+x)^2}$

(c) $L \approx 1.132$

$$1 + (y')^2 = 1 + \frac{1}{(1+x)^4}$$

$$L = \int_0^1 \sqrt{1 + \frac{1}{(1+x)^4}} dx$$

(b) $y' = 2x + 1$

$$1 + (y')^2 = 1 + 4x^2 + 4x + 1$$

$$L = \int_{-2}^1 \sqrt{2 + 4x + 4x^2} dx$$

(c) $L \approx 5.653$

4. $y = 2x^{3/2} + 3$

$$y' = 3x^{1/2}, [0, 9]$$

$$s = \int_0^9 \sqrt{1 + 9x} dx$$

$$= \left[\frac{2}{27}(1 + 9x)^{3/2} \right]_0^9$$

$$= \frac{2}{27}(82^{3/2} - 1) \approx 54.929$$

8. $y = \frac{x^5}{10} + \frac{1}{6x^3}$

$$y' = \frac{1}{2}x^4 - \frac{1}{2x^4}$$

$$1 + (y')^2 = \left(\frac{1}{2}x^4 + \frac{1}{2x^4} \right)^2, [1, 2]$$


$$s = \int_a^b \sqrt{1 + (y')^2} dx$$

$$= \int_1^2 \sqrt{\left(\frac{1}{2}x^4 + \frac{1}{2x^4} \right)^2} dx$$

$$= \int_1^2 \left(\frac{1}{2}x^4 + \frac{1}{2x^4} \right) dx$$

$$= \left[\frac{1}{10}x^5 - \frac{1}{6x^3} \right]_1^2 = \frac{779}{240} \approx 3.246$$

12. (a) $y = x^2 + x - 2, -2 \leq x \leq 1$



(b) $y' = 2x + 1$

$$1 + (y')^2 = 1 + 4x^2 + 4x + 1$$

$$L = \int_{-2}^1 \sqrt{2 + 4x + 4x^2} dx$$

(c) $L \approx 5.653$


(b) $y' = -\frac{1}{(1+x)^2}$

(c) $L \approx 1.132$

$$1 + (y')^2 = 1 + \frac{1}{(1+x)^4}$$

$$L = \int_0^1 \sqrt{1 + \frac{1}{(1+x)^4}} dx$$

16. (a) $y = \cos x, -\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$




(b) $y' = -\sin x$

$$1 + (y')^2 = 1 + \sin^2 x$$

$$L = \int_{-\pi/2}^{\pi/2} \sqrt{1 + \sin^2 x} dx$$

(c) 3.820

18. (a) $y = \ln x, 1 \leq x \leq 5$



(b) $y' = \frac{1}{x}$


$$1 + (y')^2 = 1 + \frac{1}{x^2}$$

$$L = \int_1^5 \sqrt{1 + \frac{1}{x^2}} dx$$

(c) $L \approx 4.367$

20. (a) $x = \sqrt{36 - y^2}, 0 \leq y \leq 3$

$$y = \sqrt{36 - x^2}, 3\sqrt{3} \leq x \leq 6$$



(b) $\frac{dx}{dy} = \frac{1}{2}(36 - y^2)^{-1/2}(-2y)$

$$\begin{aligned} &= \frac{-y}{\sqrt{36 - y^2}} \\ L &= \int_0^3 \sqrt{1 + \frac{y^2}{36 - y^2}} dy \\ &= \int_0^3 \frac{6}{\sqrt{36 - y^2}} dy \end{aligned}$$

(c) $L \approx 3.142 (\pi!)$

Alternatively, you can convert to a function of x .

$$y = \sqrt{36 - x^2}$$

$$y' = \frac{dy}{dx} = -\frac{x}{\sqrt{36 - x^2}}$$


$$L = \int_{3\sqrt{3}}^6 \sqrt{1 + \frac{x^2}{36 - x^2}} dx = \int_{3\sqrt{3}}^6 \frac{6}{\sqrt{36 - x^2}} dx$$

Although this integral is undefined at $x = 0$, a graphing utility still gives $L \approx 3.142$.

22. $\int_0^{\pi/4} \sqrt{1 + \left[\frac{d}{dx}(\tan x) \right]^2} dx$

$$s \approx 1$$

Matches (e)



24. $f(x) = (x^2 - 4)^2, [0, 4]$

(a) $d = \sqrt{(4 - 0)^2 + (144 - 16)^2} \approx 128.062$

$$\begin{aligned} (b) d &= \sqrt{(1 - 0)^2 + (9 - 16)^2} + \sqrt{(2 - 1)^2 + (0 - 9)^2} + \sqrt{(3 - 2)^2 + (25 - 0)^2} + \sqrt{(4 - 3)^2 + (144 - 25)^2} \\ &\approx 160.151 \end{aligned}$$

(c) $s = \int_0^4 \sqrt{1 + [4x(x^2 - 4)]^2} dx \approx 159.087$

(d) 160.287

26. Let $y = \ln x$, $1 \leq x \leq e$, $y' = \frac{1}{x}$ and $L_1 = \int_1^e \sqrt{1 + \frac{1}{x^2}} dx$.

Equivalently, $x = e^y$, $0 \leq y \leq 1$, $\frac{dx}{dy} = e^y$, and $L_2 = \int_0^1 \sqrt{1 + e^{2y}} dy = \int_0^1 \sqrt{1 + e^{2x}} dx$.

Numerically, both integrals yield $L = 2.0035$

28. $y = 31 - 10(e^{x/20} + e^{-x/20})$

$$y' = -\frac{1}{2}(e^{x/20} - e^{-x/20})$$

$$\begin{aligned} 1 + (y')^2 &= 1 + \frac{1}{4}(e^{x/10} - 2 + e^{-x/10}) = \left[\frac{1}{2}(e^{x/20} + e^{-x/20}) \right]^2 \\ s &= \int_{-20}^{20} \sqrt{\left[\frac{1}{2}(e^{x/20} + e^{-x/20}) \right]^2} dx \\ &= \frac{1}{2} \int_{-20}^{20} (e^{x/20} + e^{-x/20}) dx = \left[10(e^{x/20} - e^{-x/20}) \right]_{-20}^{20} = 20\left(e - \frac{1}{e}\right) \approx 47 \text{ ft} \end{aligned}$$

Thus, there are $100(47) = 4700$ square feet of roofing on the barn.

30. $y = 693.8597 - 68.7672 \cosh 0.0100333x$

$$y' = -0.6899619478 \sinh 0.0100333x$$

$$s = \int_{-299.2239}^{299.2239} \sqrt{1 + (-0.6899619478 \sinh 0.0100333x)^2} dx \approx 1480$$

(Use Simpson's Rule with $n = 100$ or a graphing utility.)

32. $y = \sqrt{25 - x^2}$

$$y' = \frac{-x}{\sqrt{25 - x^2}}$$

$$\begin{aligned} 1 + (y')^2 &= \frac{25}{25 - x^2} \\ s &= \int_{-3}^4 \sqrt{\frac{25}{25 - x^2}} dx \\ &= \int_{-3}^4 \frac{5}{\sqrt{25 - x^2}} dx \\ &= \left[5 \arcsin \frac{x}{5} \right]_{-3}^4 \\ &= 5 \left[\arcsin \frac{4}{5} - \arcsin \left(-\frac{3}{5} \right) \right] \approx 7.8540 \end{aligned}$$

$$\frac{1}{4}[2\pi(5)] \approx 7.8540 = s$$

34. $y = 2\sqrt{x}$

$$y' = \frac{1}{\sqrt{x}}, [4, 9]$$

$$\begin{aligned} S &= 2\pi \int_4^9 2\sqrt{x} \sqrt{1 + \frac{1}{x}} dx \\ &= 4\pi \int_4^9 \sqrt{x+1} dx \\ &= \frac{8}{3}\pi(x+1)^{3/2} \Big|_4^9 \\ &= \frac{8\pi}{3}(10^{3/2} - 5^{3/2}) \approx 171.258 \end{aligned}$$

36. $y = \frac{x}{2}$

$$y' = \frac{1}{2}$$

$$1 + (y')^2 = \frac{5}{4}, [0, 6]$$

$$\begin{aligned} S &= 2\pi \int_0^6 \frac{x}{2} \sqrt{\frac{5}{4}} dx \\ &= \left[\frac{2\pi\sqrt{5}}{8} x^2 \right]_0^6 = 9\sqrt{5} \pi \end{aligned}$$

40. $y = \ln x$

$$y' = \frac{1}{x}$$

$$1 + (y')^2 = \frac{x^2 + 1}{x^2}, [1, e]$$

$$\begin{aligned} S &= 2\pi \int_1^e x \sqrt{\frac{x^2 + 1}{x^2}} dx \\ &= 2\pi \int_1^e \sqrt{x^2 + 1} dx \approx 22.943 \end{aligned}$$

44. The surface of revolution given by f_1 will be larger. $r(x)$ is larger for f_1 .

46. $y = \sqrt{r^2 - x^2}$

$$y' = \frac{-x}{\sqrt{r^2 - x^2}}$$

$$1 + (y')^2 = \frac{r^2}{r^2 - x^2}$$


$$\begin{aligned} S &= 2\pi \int_{-r}^r \sqrt{r^2 - x^2} \sqrt{\frac{r^2}{r^2 - x^2}} dx \\ &= 2\pi \int_{-r}^r r dx = \left[2\pi rx \right]_{-r}^r = 4\pi r^2 \end{aligned}$$

42. The precalculus formula is the distance formula between two points. The representative element is

$$\sqrt{(\Delta x_i)^2 + (\Delta y_i)^2} = \sqrt{1 + \left(\frac{\Delta y_i}{\Delta x_i} \right)^2} \Delta x_i.$$

48. From Exercise 47 we have:

$$\begin{aligned} S &= 2\pi \int_0^a \frac{rx}{\sqrt{r^2 - x^2}} dx \\ &= -r\pi \int_0^a \frac{-2x dx}{\sqrt{r^2 - x^2}} \\ &= \left[-2r\pi \sqrt{r^2 - x^2} \right]_0^a \\ &= 2r^2\pi - 2r\pi\sqrt{r^2 - a^2} \\ &= 2r\pi(r - \sqrt{r^2 - a^2}) \\ &= 2\pi rh \text{ (where } h \text{ is the height of the zone)} \end{aligned}$$




50. (a) We approximate the volume by summing 6 disks of thickness 3 and circumference C_i equal to the average of the given circumferences:

$$\begin{aligned}
 V &\approx \sum_{i=1}^6 \pi r_i^2(3) = \sum_{i=1}^6 \pi \left(\frac{C_i}{2\pi}\right)^2(3) = \frac{3}{4\pi} \sum_{i=1}^6 C_i^2 \\
 &= \frac{3}{4\pi} \left[\left(\frac{50 + 65.5}{2}\right)^2 + \left(\frac{65.5 + 70}{2}\right)^2 + \left(\frac{70 + 66}{2}\right)^2 + \left(\frac{66 + 58}{2}\right)^2 + \left(\frac{58 + 51}{2}\right)^2 + \left(\frac{51 + 48}{2}\right)^2 \right] \\
 &= \frac{3}{4\pi} [57.75^2 + 67.75^2 + 68^2 + 62^2 + 54.5^2 + 49.5^2] \\
 &= \frac{3}{4\pi} [21813.625] = 5207.62 \text{ cubic inches}
 \end{aligned}$$

- (b) The lateral surface area of a frustum of a right circular cone is $\pi s(R + r)$. For the first frustum,


$$\begin{aligned}
 S_1 &\approx \pi \left[3^2 + \left(\frac{65.5 - 50}{2\pi}\right)^2 \right]^{1/2} \left[\frac{50}{2\pi} + \frac{65.5}{2\pi} \right] \\
 &= \left(\frac{50 + 65.5}{2}\right) \left[9 + \left(\frac{65.5 - 50}{2\pi}\right)^2 \right]^{1/2}.
 \end{aligned}$$



Adding the six frustums together,

$$\begin{aligned}
 S &\approx \left(\frac{50 + 65.5}{2}\right) \left[9 + \left(\frac{15.5}{2\pi}\right)^2 \right]^{1/2} + \left(\frac{65.5 + 70}{2}\right) \left[9 + \left(\frac{4.5}{2\pi}\right)^2 \right]^{1/2} + \\
 &\quad \left(\frac{70 + 66}{2}\right) \left[9 + \left(\frac{4}{2\pi}\right)^2 \right]^{1/2} + \left(\frac{66 + 58}{2}\right) \left[9 + \left(\frac{8}{2\pi}\right)^2 \right]^{1/2} + \\
 &\quad \left(\frac{58 + 51}{2}\right) \left[9 + \left(\frac{7}{2\pi}\right)^2 \right]^{1/2} + \left(\frac{51 + 48}{2}\right) \left[9 + \left(\frac{3}{2\pi}\right)^2 \right]^{1/2} \\
 &\approx 224.30 + 208.96 + 208.54 + 202.06 + 174.41 + 150.37 \\
 &= 1168.64
 \end{aligned}$$

(c) $r = 0.00401y^3 - 0.1416y^2 + 1.232y + 7.943$



(d) $V = \int_0^{18} \pi r^2 dy \approx 5275.9 \text{ cubic inches}$

$$S = \int_0^{18} 2\pi r(y) \sqrt{1 + r'(y)^2} dy$$


$$\approx 1179.5 \text{ square inches}$$

52. Individual project, see Exercise 50, 51.

54. (a) $\frac{x^2}{9} + \frac{y^2}{4} = 1$

Ellipse: $y_1 = 2\sqrt{1 - \frac{x^2}{9}}$

$y_2 = -2\sqrt{1 - \frac{x^2}{9}}$



(b) $y = 2\sqrt{1 - \frac{x^2}{9}}, 0 \leq x \leq 3$

$$y' = 2\left(\frac{1}{2}\right)\left(1 - \frac{x^2}{9}\right)^{-1/2}\left(-\frac{2x}{9}\right)$$

$$= \frac{-2x}{9\sqrt{1 - \frac{x^2}{9}}} = \frac{-2x}{3\sqrt{9 - x^2}}$$

$$L = \int_0^3 \sqrt{1 + \frac{4x^2}{81 - 9x^2}} dx$$

- (c) You cannot evaluate this definite integral, since the integrand is not defined at $x = 3$. Simpson's Rule will not work for the same reason. Also, the integrand does not have an elementary antiderivative.

56. Essay

Section 6.5 Work

2. $W = Fd = (2800)(4) = 11,200 \text{ ft} \cdot \text{lb}$

4. $W = Fd = [9(2000)]\left[\frac{1}{2}(5280)\right] = 47,520,000 \text{ ft} \cdot \text{lb}$

6. $W = \int_a^b F(x) dx$ is the work done by a force F moving an object along a straight line from $x = a$ to $x = b$.

8. (a) $W = \int_0^9 6 dx = 54 \text{ ft} \cdot \text{lbs}$

(b) $W = \int_0^7 20 dx + \int_7^9 (-10x + 90) dx = 140 + 20$
 $= 160 \text{ ft} \cdot \text{lbs}$

(c) $W = \int_0^9 \frac{1}{27} x^2 dx = \left[\frac{x^3}{81} \right]_0^9 = 9 \text{ ft} \cdot \text{lbs}$

(d) $W = \int_0^9 \sqrt{x} dx = \left[\frac{2}{3} x^{3/2} \right]_0^9 = \frac{2}{3}(27) = 18 \text{ ft} \cdot \text{lbs}$

10. $W = \int_0^{10} \frac{5}{4} x dx = \left[\frac{5}{8} x^2 \right]_6^{10}$

$= 40 \text{ in} \cdot \text{lb} \approx 3.33 \text{ ft} \cdot \text{lb}$

12. $F(x) = kx$

$800 = k(70) \Rightarrow k = \frac{80}{7}$

$W = \int_0^{70} F(x) dx = \int_0^{70} \frac{80}{7} x dx = \left[\frac{40x^2}{7} \right]_0^{70}$
 $= 28000 \text{ N} \cdot \text{cm} = 280 \text{ Nm}$

14. $F(x) = kx$

$15 = k(1) = k$

$W = 2 \int_0^4 15x dx = \left[15x^2 \right]_0^4$
 $= 240 \text{ ft} \cdot \text{lb}$

16. $W = 7.5 = \int_0^{1/6} kx dx = \left[\frac{kx^2}{2} \right]_0^{1/6} = \frac{k}{72} \Rightarrow k = 540$

$W = \int_{1/6}^{5/24} 540x dx = \left[270x^2 \right]_{1/6}^{5/24} = 4.21875 \text{ ft} \cdot \text{lbs}$

18. $W = \int_{4000}^h \frac{80,000,000}{x^2} dx = \left[-\frac{80,000,000}{x} \right]_{4000}^h$

$= \frac{-80,000,000}{h} + 20,000$

$\lim_{h \rightarrow \infty} W = 20,000 \text{ mi/ton} \approx 2.1 \times 10^{11} \text{ ft} \cdot \text{lb}$

20. Weight on surface of moon: $\frac{1}{6}(12) = 2 \text{ tons}$

Weight varies inversely as the square of distance from the center of the moon. Therefore,

$$F(x) = \frac{k}{x^2}$$

$$2 = \frac{k}{(1100)^2}$$

$$k = 2.42 \times 10^6$$

$$W = \int_{1100}^{1150} \frac{2.42 \times 10^6}{x^2} dx = \left[\frac{-2.42 \times 10^6}{x} \right]_{1100}^{1150} = 2.42 \times 10^6 \left(\frac{1}{1100} - \frac{1}{1150} \right)$$

$$\approx 95.652 \text{ mi} \cdot \text{ton} \approx 1.01 \times 10^9 \text{ ft} \cdot \text{lb}$$

22. The bottom half had to be pumped a greater distance than the top half.

24. Volume of disk: $4\pi \Delta y$

Weight of disk: $9800(4\pi) \Delta y$

Distance the disk of water is moved: y

$$\begin{aligned} W &= \int_{10}^{12} y(9800)(4\pi) dy = 39,200\pi \left[\frac{y^2}{2} \right]_{10}^{12} \\ &= 39,200\pi(22) \\ &= 862,400\pi \text{ newton-meters} \end{aligned}$$

26. Volume of disk: $\pi \left(\frac{2}{3}y \right)^2 \Delta y$

Weight of disk: $62.4\pi \left(\frac{2}{3}y \right)^2 \Delta y$

Distance: y


$$\begin{aligned} (a) \quad W &= \frac{4}{9}(62.4)\pi \int_0^2 y^3 dy = \left[\frac{4}{9}(62.4)\pi \left(\frac{1}{4}y^4 \right) \right]_0^2 \approx 110.9\pi \text{ ft} \cdot \text{lb} \\ (b) \quad W &= \frac{4}{9}(62.4)\pi \int_4^6 y^3 dy = \left[\frac{4}{9}(62.4)\pi \left(\frac{1}{4}y^4 \right) \right]_4^6 \approx 7210.7\pi \text{ ft} \cdot \text{lb} \end{aligned}$$

28. Volume of each layer: $\frac{y+3}{3}(3) \Delta y = (y+3) \Delta y$

Weight of each layer: $55.6(y+3) \Delta y$

Distance: $6 - y$

$$\begin{aligned} W &= \int_0^3 55.6(6-y)(y+3) dy = 55.6 \int_0^3 (18 + 3y - y^2) dy \\ &= 55.6 \left[18y + \frac{3y^2}{2} - \frac{y^3}{3} \right]_0^3 \\ &= 3252.6 \text{ ft} \cdot \text{lb} \end{aligned}$$




30. Volume of layer: $V = 12(2)\sqrt{(25/4) - y^2} \Delta y$

Weight of layer: $W = 42(24)\sqrt{(25/4) - y^2} \Delta y$


Distance: $\frac{19}{2} - y$

$$\begin{aligned} W &= \int_{-2.5}^{2.5} 42(24) \sqrt{\frac{25}{4} - y^2} \left(\frac{19}{2} - y \right) dy \\ &= 1008 \left[\frac{19}{2} \int_{-2.5}^{2.5} \sqrt{\frac{25}{4} - y^2} dy + \int_{-2.5}^{2.5} \sqrt{\frac{25}{4} - y^2} (-y) dy \right] \end{aligned}$$



The second integral is zero since the integrand is odd and the limits of integration are symmetric to the origin. The first integral represents the area of a semicircle of radius $\frac{5}{2}$. Thus, the work is

$$W = 1008 \left(\frac{19}{2} \right) \pi \left(\frac{5}{2} \right)^2 \left(\frac{1}{2} \right) = 29,925\pi \text{ ft} \cdot \text{lb} \approx 94,012.16 \text{ ft} \cdot \text{lb}.$$



32. The lower 10 feet of chain are raised 5 feet with a constant force.

$$W_1 = 3(10)5 = 150 \text{ ft} \cdot \text{lb}$$

The top 5 feet will be raised with variable force.

Weight of section: $3 \Delta y$

Distance: $5 - y$

$$W_2 = 3 \int_0^5 (5 - y) dy = \left[-\frac{3}{2}(5 - y)^2 \right]_0^5 = \frac{75}{2} \text{ ft} \cdot \text{lb}$$

$$W = W_1 + W_2 = 150 + \frac{75}{2} = \frac{375}{2} \text{ ft} \cdot \text{lb}$$

34. The work required to lift the chain is 337.5 ft · lb (from Exercise 31). The work required to lift the 500-pound load is $W = (500)(15) = 7500$. The work required to lift the chain with a 100-pound load attached is

$$W = 337.5 + 7500 = 7837.5 \text{ ft} \cdot \text{lbs}$$

36. $W = 3 \int_0^6 (12 - 2y) dy = \left[-\frac{3}{4}(12 - 2y)^2 \right]_0^6 = \frac{3}{4}(12)^2 = 108 \text{ ft} \cdot \text{lb}$

38. Work to pull up the ball: $W_1 = 500(40) = 20,000 \text{ ft} \cdot \text{lb}$

$$40. \quad p = \frac{k}{V}$$

Work to pull up the cable: force is variable

Weight per section: $1 \Delta y$

Distance: $40 - x$

$$W_2 = \int_0^{40} (40 - x) dx = \left[-\frac{1}{2}(40 - x)^2 \right]_0^{40} = 800 \text{ ft} \cdot \text{lb}$$

$$W = W_1 + W_2 = 20,000 + 800 = 20,800 \text{ ft} \cdot \text{lb}$$

$$2500 = \frac{k}{1} \Rightarrow k = 2500$$


$$W = \int_1^3 \frac{2500}{V} dV = \left[2500 \ln V \right]_1^3 = 2500 \ln 3 \approx 2746.53 \text{ ft} \cdot \text{lb}$$

42. (a) $W = FD = (8000\pi)(2) = 16,000\pi \text{ ft} \cdot \text{lbs}$

$$(b) W \approx \frac{2 - 0}{3(6)} [0 + 4(20,000) + 2(22,000) + 4(15,000) + 2(10,000) + 4(5000) + 0]$$

$$\approx 24,88.889 \text{ ft} \cdot \text{lb}$$

$$(c) F(x) = -16,261.36x^4 + 85,295.45x^3 - 157,738.64x^2 + 104,386.36x - 32.4675$$



- (d) $F(x) = 0$ when $x \approx 0.524$ feet. $F(x)$ is a maximum when $x \approx 0.524$ feet.

$$(e) W = \int_0^2 F(x) dx \approx 25,180.5 \text{ ft} \cdot \text{lbs}$$

44. $W = \int_0^4 \left(\frac{e^{x^2} - 1}{100} \right) dx \approx 11,494 \text{ ft} \cdot \text{lb}$

46. $W = \int_0^2 1000 \sinh x dx \approx 2762.2 \text{ ft} \cdot \text{lb}$

Section 6.6 Moments, Centers of Mass, and Centroids

2. $\bar{x} = \frac{7(-3) + 4(-2) + 3(5) + 8(6)}{7 + 4 + 3 + 8} = \frac{17}{11}$

4. $\bar{x} = \frac{12(-6) + 1(-4) + 6(-2) + 3(0) + 11(8)}{12 + 1 + 6 + 3 + 11} = 0$

6. The center of mass is translated k units as well.

8. $200x = 550(5 - x)$ (Person on left)

$$200x = 2750 - 550x$$


$$750x = 2750$$

$$x = 3\frac{2}{3} \text{ feet}$$

10. $\bar{x} = \frac{10(1) + 2(5) + 5(-4)}{10 + 2 + 5} = 0$

$$\bar{y} = \frac{10(-1) + 2(5) + 5(0)}{10 + 2 + 5} = 0$$


$$(\bar{x}, \bar{y}) = (0, 0)$$



12. $\bar{x} = \frac{12(2) + 6(-1) + \frac{15}{2}(6) + 15(2)}{12 + 6 + \frac{15}{2} + 15} = \frac{93}{40.5} = \frac{62}{27}$

$$\bar{y} = \frac{12(3) + 6(5) + \frac{15}{2}(8) + 15(-2)}{12 + 6 + \frac{15}{2} + 15} = \frac{96}{40.5} = \frac{64}{27}$$

$$(\bar{x}, \bar{y}) = \left(\frac{62}{27}, \frac{64}{27}\right)$$



14. $m = \rho \int_0^2 \frac{1}{2}x^2 dx = \left[\rho \frac{x^3}{6} \right]_0^2 = \frac{4}{3}\rho$


$$M_x = \rho \int_0^2 \frac{1}{2} \left(\frac{1}{2}x^2 \right) \left(\frac{1}{2}x^2 \right) dx = \frac{\rho}{8} \int_0^2 x^4 dx = \left[\frac{\rho}{40}x^5 \right]_0^2 = \frac{32}{40}\rho = \frac{4}{5}\rho$$

$$\bar{y} = \frac{M_x}{m} = \frac{\frac{4}{5}\rho}{\frac{4}{3}\rho} = \frac{3}{5}$$


$$M_y = \rho \int_0^2 x \left(\frac{1}{2}x^2 \right) dx = \frac{1}{2}\rho \int_0^2 x^3 dx = \left[\frac{\rho}{8}x^4 \right]_0^2 = 2\rho$$

$$\bar{x} = \frac{M_y}{m} = \frac{2\rho}{\frac{4}{3}\rho} = \frac{3}{2}$$


$$(\bar{x}, \bar{y}) = \left(\frac{3}{2}, \frac{3}{5}\right)$$



$$\begin{aligned}
 16. \quad m &= \rho \int_0^1 (\sqrt{x} - x) dx = \rho \left[\frac{2}{3}x^{3/2} - \frac{x^2}{2} \right]_0^1 = \frac{\rho}{6} \\
 M_x &= \rho \int_0^1 \frac{(\sqrt{x} + x)}{2} (\sqrt{x} - x) dx = \frac{\rho}{2} \int_0^1 (x - x^2) dx = \frac{\rho}{2} \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = \frac{\rho}{12} \\
 \bar{y} &= \frac{M_x}{m} = \frac{\rho}{12} \left(\frac{6}{\rho} \right) = \frac{1}{2} \\
 M_y &= \rho \int_0^1 x(\sqrt{x} - x) dx = \rho \int_0^1 (x^{3/2} - x^2) dx = \rho \left[\frac{2}{5}x^{5/2} - \frac{x^3}{3} \right]_0^1 = \frac{\rho}{15} \\
 \bar{x} &= \frac{M_y}{m} = \frac{\rho}{15} \left(\frac{6}{\rho} \right) = \frac{2}{5} \\
 (\bar{x}, \bar{y}) &= \left(\frac{2}{5}, \frac{1}{2} \right)
 \end{aligned}$$



$$\begin{aligned}
 18. \quad m &= \rho \int_0^9 \left[(\sqrt{x} + 1) - \left(\frac{1}{3}x + 1 \right) \right] dx = \rho \int_0^9 \left(\sqrt{x} - \frac{1}{3}x \right) dx \\
 &= \rho \left[\frac{2}{3}x^{3/2} - \frac{x^2}{6} \right]_0^9 = \rho \left(18 - \frac{27}{2} \right) = \frac{9}{2}\rho \\
 M_x &= \rho \int_0^9 \frac{\sqrt{x} + 1 + \frac{1}{3}x + 1}{2} \left(\sqrt{x} + 1 - \frac{1}{3}x - 1 \right) dx = \frac{\rho}{2} \int_0^9 \left(\sqrt{x} + \frac{1}{3}x + 2 \right) \left(\sqrt{x} - \frac{1}{3}x \right) dx \\
 &= \frac{\rho}{2} \int_0^9 \left(x - \frac{1}{3}x^{3/2} + \frac{1}{3}x^{3/2} - \frac{1}{9}x^2 + 2\sqrt{x} - \frac{2}{3}x \right) dx = \frac{\rho}{2} \int_0^9 \left(\frac{1}{3}x - \frac{1}{9}x^2 + 2\sqrt{x} \right) dx \\
 &= \frac{\rho}{2} \left[\frac{x^2}{6} - \frac{x^3}{27} + \frac{4}{3}x^{3/2} \right]_0^9 = \frac{\rho}{2} \left[\frac{27}{2} - 27 + 36 \right] = \frac{45}{3}\rho \\
 M_y &= \rho \int_0^9 x \left[\sqrt{x} + 1 - \frac{1}{3}x - 1 \right] dx = \rho \int_0^9 \left(x^{3/2} - \frac{1}{3}x^2 \right) dx = \rho \left[\frac{2}{5}x^{5/2} - \frac{1}{9}x^3 \right]_0^9 \\
 &= \rho \left[\frac{486}{5} - 81 \right] = \frac{81}{5}\rho \\
 \bar{x} &= \frac{M_y}{m} = \frac{\frac{81}{5}\rho}{\frac{45}{3}\rho} = \frac{18}{5}; \bar{y} = \frac{M_x}{m} = \frac{\frac{45}{3}\rho}{\frac{45}{3}\rho} = \frac{5}{2} \\
 (\bar{x}, \bar{y}) &= \left(\frac{18}{5}, \frac{5}{2} \right)
 \end{aligned}$$




$$20. \quad m = 2\rho \int_0^8 (4 - x^{2/3}) dx = 2\rho \left[4x - \frac{3}{5}x^{5/3} \right]_0^8 = \frac{128\rho}{5}$$

By symmetry, M_y and $\bar{x} = 0$.

$$\begin{aligned} M_x &= 2\rho \int_0^8 \left(\frac{4 + x^{2/3}}{2} \right) (4 - x^{2/3}) dx = \rho \left[16x - \frac{3}{7}x^{7/3} \right]_0^8 = \frac{512\rho}{7} \\ \bar{y} &= \frac{512\rho}{7} \left(\frac{5}{128\rho} \right) = \frac{20}{7} \end{aligned}$$

$$(\bar{x}, \bar{y}) = \left(0, \frac{20}{7} \right)$$



$$22. \quad m = \rho \int_0^2 (2y - y^2) dy = \rho \left[y^2 - \frac{y^3}{3} \right]_0^2 = \frac{4\rho}{3}$$


$$M_y = \rho \int_0^2 \left(\frac{2y - y^2}{2} \right) (2y - y^2) dy = \frac{\rho}{2} \left[\frac{4y^3}{3} - y^4 + \frac{y^5}{5} \right]_0^2 = \frac{8\rho}{15}$$

$$\bar{x} = \frac{M_y}{m} = \frac{8\rho}{15} \left(\frac{3}{4\rho} \right) = \frac{2}{5}$$

$$M_x = \rho \int_0^2 y(2y - y^2) dy = \rho \left[\frac{2y^3}{3} - \frac{y^4}{4} \right]_0^2 = \frac{4\rho}{3}$$

$$\bar{y} = \frac{M_x}{m} = \frac{4\rho}{3} \left(\frac{3}{4\rho} \right) = 1$$

$$(\bar{x}, \bar{y}) = \left(\frac{2}{5}, 1 \right)$$



$$24. \quad m = \rho \int_{-1}^2 [(y+2) - y^2] dy = \rho \left[\frac{y^2}{2} + 2y - \frac{y^3}{3} \right]_{-1}^2 = \frac{9\rho}{2}$$


$$\begin{aligned} M_y &= \rho \int_{-1}^2 \frac{[(y+2) + y^2]}{2} [(y+2) - y^2] dy \\ &= \frac{\rho}{2} \int_{-1}^2 [(y+2)^2 - y^4] dy = \frac{\rho}{2} \left[\frac{(y+2)^3}{3} - \frac{y^5}{5} \right]_{-1}^2 = \frac{36\rho}{5} \end{aligned}$$

$$\bar{x} = \frac{M_y}{m} = \frac{36\rho}{5} \left(\frac{2}{9\rho} \right) = \frac{8}{5}$$

$$\begin{aligned} M_x &= \rho \int_{-1}^2 y[(y+2) - y^2] dy \\ &= \rho \int_{-1}^2 (2y + y^2 - y^3) dy = \rho \left[y^2 + \frac{y^3}{3} - \frac{y^4}{4} \right]_{-1}^2 = \frac{9\rho}{4} \end{aligned}$$

$$\bar{y} = \frac{M_x}{m} = \frac{9\rho}{4} \left(\frac{2}{9\rho} \right) = \frac{1}{2}$$

$$(\bar{x}, \bar{y}) = \left(\frac{8}{5}, \frac{1}{2} \right)$$



$$26. \quad A = \int_1^4 \frac{1}{x} dx = \left[\ln|x| \right]_1^4 = \ln 4$$

$$M_x = \frac{1}{2} \int_1^4 \frac{1}{x^2} dx = \left[\frac{1}{2} \left(-\frac{1}{x} \right) \right]_1^4 = \left(-\frac{1}{8} + \frac{1}{2} \right) = \frac{3}{8}$$

$$M_y = \int_1^4 x \left(\frac{1}{x} \right) dx = \left[x \right]_1^4 = 3$$

$$28. A = \int_{-2}^2 -(x^2 - 4) dx = 2 \int_0^2 (4 - x^2) dx = \left[8x - \frac{2x^3}{3} \right]_0^2 = 16 - \frac{16}{3} = \frac{32}{3}$$

$$\begin{aligned} M_x &= \frac{1}{2} \int_{-2}^2 (x^2 - 4)(4 - x^2) dx = -\frac{1}{2} \int_{-2}^2 (x^4 - 8x^2 + 16) dx \\ &= -\frac{1}{2} \left[\frac{x^5}{5} - \frac{8x^3}{3} + 16x \right]_{-2}^2 = -\left[\frac{32}{5} - \frac{64}{3} + 32 \right] = -\frac{256}{15} \end{aligned}$$

$M_y = 0$ by symmetry.

$$30. m = \rho \int_0^4 xe^{-x/2} dx \approx 2.3760\rho$$


$$M_x = \rho \int_0^4 \left(\frac{xe^{-x/2}}{2} \right) (xe^{-x/2}) dx = \frac{\rho}{2} \int_0^4 x^2 e^{-x} dx \approx 0.7619\rho$$

$$M_y = \rho \int_0^4 x^2 e^{-x/2} dx \approx 5.1732\rho$$

$$\bar{x} = \frac{M_y}{m} \approx 2.2$$

$$\bar{y} = \frac{M_x}{m} \approx 0.3$$

Therefore, the centroid is (2.2, 0.3).




$$32. m = \rho \int_{-2}^2 \frac{8}{x^2 + 4} dx \approx 6.2832\rho$$

$$M_x = \rho \int_{-2}^2 \frac{1}{2} \left(\frac{8}{x^2 + 4} \right) \left(\frac{8}{x^2 + 4} \right) dx = 32\rho \int_{-2}^2 \frac{1}{(x^2 + 4)^2} dx \approx 5.14149\rho$$

$$\bar{y} = \frac{M_x}{m} \approx 0.8$$

$\bar{x} = 0$ by symmetry. Therefore, the centroid is (0, 0.8).




$$34. A = bh = ac$$

$$\frac{1}{A} = \frac{1}{ac}$$

$$\begin{aligned} \bar{x} &= \frac{1}{ac} \frac{1}{2} \int_0^c \left[\left(\frac{b}{c}y + a \right)^2 - \left(\frac{b}{c}y \right)^2 \right] dy \\ &= \frac{1}{2ac} \int_0^c \left(\frac{2ab}{c}y + a^2 \right) dy \\ &= \frac{1}{2ac} \left[\frac{ab}{c}y^2 + a^2y \right]_0^c = \frac{1}{2ac} [abc + a^2c] = \frac{1}{2}(b + a) \end{aligned}$$

$$\bar{y} = \frac{1}{ac} \int_0^c y \left[\left(\frac{b}{c}y + a \right) - \left(\frac{b}{c}y \right) \right] dy = \left[\frac{1}{c} \frac{y^2}{2} \right]_0^c = \frac{c}{2}$$

$$(\bar{x}, \bar{y}) = \left(\frac{b+a}{2}, \frac{c}{2} \right)$$



This is the point of intersection of the diagonals.

36. $\bar{x} = 0$ by symmetry


$$A = \frac{1}{2}\pi r^2$$

$$\frac{1}{A} = \frac{2}{\pi r^2}$$

$$\bar{y} = \frac{2}{\pi r^2} \frac{1}{2} \int_{-r}^r (\sqrt{r^2 - x^2})^2 dx$$

$$= \frac{1}{\pi r^2} \left[r^2 x - \frac{x^3}{3} \right]_{-r}^r = \frac{1}{\pi r^2} \left[\frac{4r^3}{3} \right] = \frac{4r}{3\pi}$$

$$(\bar{x}, \bar{y}) = \left(0, \frac{4r}{3\pi}\right)$$



38. $A = \int_0^1 [1 - (2x - x^2)] dx = \frac{1}{3}$

$$\frac{1}{A} = 3$$

$$\bar{x} = 3 \int_0^1 x[1 - (2x - x^2)] dx = 3 \int_0^1 [x - 2x^2 + x^3] dx = 3 \left[\frac{x^2}{2} - \frac{2}{3}x^3 + \frac{x^4}{4} \right]_0^1 = \frac{1}{4}$$

$$\bar{y} = 3 \int_0^1 \frac{[1 + (2x - x^2)]}{2} [1 - (2x - x^2)] dx = \frac{3}{2} \int_0^1 [1 - (2x - x^2)^2] dx$$

$$= \frac{3}{2} \int_0^1 [1 - 4x^2 + 4x^3 - x^4] dx = \frac{3}{2} \left[x - \frac{4}{3}x^3 + x^4 - \frac{x^5}{5} \right]_0^1 = \frac{7}{10}$$

$$(\bar{x}, \bar{y}) = \left(\frac{1}{4}, \frac{7}{10}\right)$$

40. (a) $M_y = 0$ by symmetry

$$M_y = \int_{-\sqrt[2n]{b}}^{\sqrt[2n]{b}} x(b - x^{2n}) dx = 0$$

because $bx - x^{2n+1}$ is an odd function.

(c) $M_x = \int_{-\sqrt[2n]{b}}^{\sqrt[2n]{b}} \frac{(b + x^{2n})(b - x^{2n})}{2} dx = \int_{-\sqrt[2n]{b}}^{\sqrt[2n]{b}} \frac{1}{2}(b^2 - x^{4n}) dx$

$$= \frac{1}{2} \left(b^2 x - \frac{x^{4n+1}}{4n+1} \right) \Big|_{-\sqrt[2n]{b}}^{\sqrt[2n]{b}}$$

$$= b^2 b^{1/2n} - \frac{b^{(4n+1)/2n}}{4n+1} = \frac{4n}{4n+1} b^{(4n+1)/2n}$$

$$A = \int_{-\sqrt[2n]{b}}^{\sqrt[2n]{b}} (b - x^{2n}) dx = 2 \left[bx - \frac{x^{2n+1}}{2n+1} \right]_0^{\sqrt[2n]{b}}$$

$$= 2 \left[b \cdot b^{1/2n} - \frac{b^{(2n+1)/2n}}{2n+1} \right] = \frac{4n}{2n+1} b^{(2n+1)/2n}$$

$$\bar{y} = \frac{M_x}{A} = \frac{4n b^{(4n+1)/2n}/(4n+1)}{4n b^{(2n+1)/2n}/(2n+1)} = \frac{2n+1}{4n+1} b$$


(b) $\bar{y} > \frac{b}{2}$ because there is more area above $y = \frac{b}{2}$ than below.

(d)

n	1	2	3	4
\bar{y}	$\frac{3}{5}b$	$\frac{5}{9}b$	$\frac{7}{13}b$	$\frac{9}{17}b$

(e) $\lim_{n \rightarrow \infty} \bar{y} = \lim_{n \rightarrow \infty} \frac{2n+1}{4n+1} b = \frac{1}{2} b$

(f) As $n \rightarrow \infty$, the figure gets narrower.



42. Let $f(x)$ be the top curve, given by $l + d$. The bottom curve is $d(x)$.

x	0	0.5	1.0	1.5	2.0
f	2.0	1.93	1.73	1.32	0
d	0.50	0.48	0.43	0.33	0

$$\begin{aligned}
 \text{(a) Area} &= 2 \int_0^2 [f(x) - d(x)] dx \\
 &\approx 2 \frac{2}{3(4)} [1.50 + 4(1.45) + 2(1.30) + 4(.99) + 0] \\
 &= \frac{1}{3}[13.86] = 4.62 \\
 M_x &= \int_{-2}^2 \frac{f(x) + d(x)}{2} (f(x) - d(x)) dx \\
 &= \int_0^2 [f(x)^2 - d(x)^2] dx \\
 &= \frac{2}{3(4)} [3.75 + 4(3.4945) + 2(2.808) + 4(1.6335) + 0] \\
 &= \frac{1}{6}[29.878] = 4.9797 \\
 \bar{y} &= \frac{M_x}{A} = \frac{4.9797}{4.62} = 1.078
 \end{aligned}$$

$$(\bar{x}, \bar{y}) = (0, 1.078)$$

44. Centroids of the given regions: $\left(\frac{1}{2}, \frac{3}{2}\right)$, $\left(2, \frac{1}{2}\right)$, and $\left(\frac{7}{2}, 1\right)$

Area: $A = 3 + 2 + 2 = 7$

$$\begin{aligned}
 \bar{x} &= \frac{3(1/2) + 2(2) + 2(7/2)}{7} = \frac{25/2}{7} = \frac{25}{14} \\
 \bar{y} &= \frac{3(3/2) + 2(1/2) + 2(1)}{7} = \frac{15/2}{7} = \frac{15}{14}
 \end{aligned}$$

$$(\bar{x}, \bar{y}) = \left(\frac{25}{14}, \frac{15}{14}\right)$$

46. $m_1 = \frac{7}{8}(2) = \frac{7}{4}$, $P_1 = \left(0, \frac{7}{16}\right)$




$$m_2 = \frac{7}{8}\left(6 - \frac{7}{8}\right) = \frac{287}{64}, P_2 = \left(0, \frac{55}{16}\right)$$

By symmetry, $\bar{x} = 0$.

$$\bar{y} = \frac{(7/4)(7/16) + (287/64)(55/16)}{(7/4) + (287/64)} = \frac{16.569}{6384} = \frac{5523}{2128}$$

$$(\bar{x}, \bar{y}) = \left(0, \frac{5523}{2128}\right) \approx (0, 2.595)$$

$$\begin{aligned}
 \text{(b) } f(x) &= -0.1061x^4 - 0.06126x^2 + 1.9527 \\
 d(x) &= -0.02648x^4 - 0.01497x^2 + .4862 \\
 \text{(c) } \bar{y} &= \frac{M_x}{A} \approx \frac{4.9133}{4.59998} = 1.068 \\
 (\bar{x}, \bar{y}) &= (0, 1.068)
 \end{aligned}$$



48. Centroids of the given regions: $(3, 0)$ and $(1, 0)$

Mass: $8 + \pi$

$$\bar{y} = 0$$

$$\bar{x} = \frac{8(1) + \pi(3)}{8 + \pi} = \frac{8 + 3\pi}{8 + \pi}$$

$$(\bar{x}, \bar{y}) = \left(\frac{8 + 3\pi}{8 + \pi}, 0 \right) \approx (1.56, 0)$$

52. $A = \int_2^6 2\sqrt{x-2} dx = \frac{4}{3}(x-2)^{3/2} \Big|_2^6 = \frac{32}{3}$

$$M_y = \int_2^6 (x)2\sqrt{x-2} dx = 2 \int_2^6 x\sqrt{x-2} dx$$

Let $u = x - 2, x = u + 2, du = dx$:

$$\begin{aligned} M_y &= 2 \int_0^4 (u+2)\sqrt{u} du = 2 \int_0^4 (u^{3/2} + 2u^{1/2}) du = 2 \left[\frac{2}{5}u^{5/2} + \frac{4}{3}u^{3/2} \right]_0^4 \\ &= 2 \left[\frac{64}{5} + \frac{32}{3} \right] = \frac{704}{15} \end{aligned}$$

$$\bar{x} = \frac{M_y}{A} = \frac{704/15}{32/3} = \frac{22}{5}$$

$$r = \bar{x} = \frac{22}{5}$$

$$V = 2\pi r A = 2\pi \left(\frac{22}{5} \right) \left(\frac{32}{3} \right) = \frac{1408\pi}{15} \approx 294.89$$



54. A planar lamina is a thin flat plate of constant density. The center of mass (\bar{x}, \bar{y}) is the balancing point on the lamina.

56. Let R be a region in a plane and let L be a line such that L does not intersect the interior of R . If r is the distance between the centroid of R and L , then the volume V of the solid of revolution formed by revolving R about L is

$$V = 2\pi r A$$

where A is the area of R .

58. The centroid of the circle is $(1, 0)$. The distance traveled by the centroid is 2π . The arc length of the circle is also 2π . Therefore, $S = (2\pi)(2\pi) = 4\pi^2$.



50. $V = 2\pi r A = 2\pi(3)(4\pi) = 24\pi^2$

Section 6.7 Fluid Pressure and Fluid Force

2. $F = PA = [62.4(5)](16) = 4992 \text{ lb}$


4. $F = 62.4(h + 4)(48) - (62.4)(h)(48)$

$$= 62.4(4)(48) = 11,980.8 \text{ lb}$$

6. $h(y) = 3 - y$


$$\begin{aligned} L(y) &= \frac{4}{3}y \\ F &= 62.4 \int_0^3 (3-y)\left(\frac{4}{3}y\right) dy \\ &= \frac{4}{3}(62.4) \int_0^3 (3y - y^2) dy \\ &= \frac{4}{3}(62.4) \left[\frac{3y^2}{2} - \frac{y^3}{3} \right]_0^3 = 374.4 \text{ lb} \end{aligned}$$

Force is one-third that of Exercise 5.




8. $h(y) = -y$

$$\begin{aligned} L(y) &= 2\sqrt{4 - y^2} \\ F &= 62.4 \int_{-2}^0 (-y)(2)\sqrt{4 - y^2} dy \\ &= \left[62.4 \left(\frac{2}{3} \right) (4 - y^2)^{3/2} \right]_{-2}^0 = 332.8 \text{ lb} \end{aligned}$$



10. $h(y) = -y$

$$\begin{aligned} L(y) &= \frac{4}{3}\sqrt{9 - y^2} \\ F &= 62.4 \int_{-3}^0 (-y)\frac{4}{3}\sqrt{9 - y^2} dy \\ &= 62.4 \left(\frac{2}{3} \right) \int_{-3}^0 (9 - y^2)^{1/2}(-2y) dy \\ &= \left[62.4 \left(\frac{4}{9} \right) (9 - y^2)^{3/2} \right]_{-3}^0 = 748.8 \text{ lb} \end{aligned}$$




12. $h(y) = (1 + 3\sqrt{2}) - y$

$L_1(y) = 2y$ (lower part)


$L_2(y) = 2(3\sqrt{2} - y)$ (upper part)

$$\begin{aligned} F &= 2(9800) \left[\int_0^{3\sqrt{2}/2} (1 + 3\sqrt{2} - y)y dy + \int_{3\sqrt{2}/2}^{3\sqrt{2}} (1 + 3\sqrt{2} - y)(3\sqrt{2} - y) dy \right] \\ &= 19,600 \left[\frac{y^2}{2} - 3\sqrt{2}y - \frac{y^3}{3} \right]_0^{3\sqrt{2}/2} + \left[3\sqrt{2}y + 18y + \frac{y^3}{3} - \frac{6\sqrt{2} + 1}{2}y \right]_{3\sqrt{2}/2}^{3\sqrt{2}} \\ &= 19,600 \left[\frac{9(2\sqrt{2} + 1)}{4} + \frac{9(\sqrt{2} + 1)}{4} \right] \\ &= 44,100(3\sqrt{2} + 2) \text{ Newtons} \end{aligned}$$




14. $h(y) = 6 - y$

$$\begin{aligned} L(y) &= 1 \\ F &= 9800 \int_0^5 1(6 - y) dy \\ &= 9800 \left[6y - \frac{y^2}{2} \right]_0^5 = 171,500 \text{ Newtons} \end{aligned}$$




16. $h(y) = -y$

$$\begin{aligned} L(y) &= 2 \left(\frac{4}{3} \sqrt{9 - y^2} \right) \\ F &= 140.7 \int_{-3}^0 (-y)(2) \left(\frac{4}{3} \sqrt{9 - y^2} \right) dy \\ &= \frac{(140.7)(4)}{3} \int_{-3}^0 \sqrt{9 - y^2} (-2y) dy \\ &= \left[\frac{(140.7)(4)}{3} \left(\frac{2}{3} \right) (9 - y^2)^{3/2} \right]_{-3}^0 = 3376.8 \text{ lb} \end{aligned}$$



18. $h(y) = -y$

$$\begin{aligned} L(y) &= 5 + \frac{5}{3}y \\ F &= 140.7 \int_{-3}^0 (-y) \left(5 + \frac{5}{3}y \right) dy \\ &= 140.7 \int_{-3}^0 \left(-5y - \frac{5}{3}y^2 \right) dy \\ &= 140.7 \left[-\frac{5}{2}y^2 - \frac{5}{9}y^3 \right]_{-3}^0 \\ &= 140.7 \left[\frac{45}{2} - 15 \right] = 1055.25 \text{ lb} \end{aligned}$$



20. $h(y) = \frac{3}{2} - y$

$$\begin{aligned} L(y) &= 2 \left(\frac{1}{2} \right) \sqrt{9 - 4y^2} \\ F &= 42 \int_{-3/2}^{3/2} \left(\frac{3}{2} - y \right) \sqrt{9 - 4y^2} dy = 63 \int_{-3/2}^{3/2} \sqrt{9 - 4y^2} dy + \frac{21}{4} \int_{-3/2}^{3/2} \sqrt{9 - 4y^2} (-8y) dy \end{aligned}$$

The second integral is zero since it is an odd function and the limits of integration are symmetric to the origin. The first integral is twice the area of a semicircle of radius $\frac{3}{2}$.

$$(\sqrt{9 - 4y^2} = 2\sqrt{(9/4) - y^2})$$

Thus, the force is $63(\frac{9}{4}\pi) = 141.75\pi \approx 445.32 \text{ lb.}$

22. (a) $F = wk\pi r^2 = (62.4)(7)(\pi 2^2) = 1747.2\pi \text{ lbs}$

(b) $F = wk\pi r^2 = (62.4)(5)(\pi 3^2) = 2808\pi \text{ lbs}$

24. (a) $F = wkhb = (62.4)\left(\frac{11}{2}\right)(3)(5) = 5148 \text{ lbs}$

(b) $F = wkhb = (62.4)\left(\frac{17}{5}\right)(5)(10) = 10,608 \text{ lbs}$

- 26.** From Exercise 21:

$$F = 64(15)\pi\left(\frac{1}{2}\right)^2 \approx 753.98 \text{ lb}$$

- 28.** $h(y) = 3 - y$


Solving $y = 5x^2/(x^2 + 4)$ for x , you obtain

$$x = \sqrt{4y/(5 - y)}.$$

$$L(y) = 2 \sqrt{\frac{4y}{5-y}}$$

$$F = 62.4(2) \int_0^3 (3 - y) \sqrt{\frac{4y}{5 - y}} dy$$

$$= 2(124.8) \int_0^3 (3 - y) \sqrt{\frac{y}{5 - y}} dy \approx 546.265 \text{ lb}$$




- 32.** Fluid pressure is the force per unit of area exerted by a fluid over the surface of a body.

- 30.** $h(y) = 12 - y$

$$L(y) = 2 \frac{\sqrt{7(16 - y^2)}}{2} = \sqrt{7(16 - y^2)}$$

$$F = 62.4 \int_0^4 (12 - y) \sqrt{7(16 - y^2)} \, dy$$

$$= 62.4\sqrt{7} \int_0^4 (12 - y)\sqrt{16 - y^2} dy \approx 21373.7 \text{ lb}$$




- 34.** The left window experiences the greater fluid force because its centroid is lower.


Review Exercises for Chapter 6

$$2. A = \int_{1/2}^5 \left(4 - \frac{1}{x^2}\right) dx$$


$$= \left[4x + \frac{1}{x}\right]_{1/2}^5 = \frac{81}{5}$$




$$\begin{aligned}
 4. A &= \int_0^1 [(y^2 - 2y) - (-1)] dy \\
 &= \int_0^1 (y^2 - 2y + 1) dy \\
 &= \int_0^1 (y - 1)^2 dy \\
 &= \left[\frac{(y - 1)^3}{3} \right]_0^1 = \frac{1}{3}
 \end{aligned}$$




$$\begin{aligned}
 6. A &= \int_{-1}^2 [(y+3) - (y^2 + 1)] dy \\
 &= \int_{-1}^2 (2 + y - y^2) dy \\
 &= \left[2y + \frac{1}{2}y^2 - \frac{1}{3}y^3 \right]_{-1}^2 = \frac{9}{2}
 \end{aligned}$$




$$\begin{aligned}
 8. A &= 2 \int_{\pi/6}^{\pi/2} (2 - \csc x) dx \\
 &= 2 \left[2x - \ln|\csc x - \cot x| \right]_{\pi/6}^{\pi/2} \\
 &= 2 \left([\pi - 0] - \left[\frac{\pi}{3} - \ln(2 - \sqrt{3}) \right] \right) \\
 &= 2 \left[\frac{2\pi}{3} + \ln(2 - \sqrt{3}) \right] \approx 1.555
 \end{aligned}$$



$$\begin{aligned}
 10. A &= \int_{\pi/3}^{5\pi/3} \left(\frac{1}{2} - \cos y \right) dy + \int_{5\pi/3}^{7\pi/3} \left(\cos y - \frac{1}{2} \right) dy \\
 &= \left[\frac{y}{2} - \sin y \right]_{\pi/3}^{5\pi/3} + \left[\sin y - \frac{y}{2} \right]_{5\pi/3}^{7\pi/3} \\
 &= \frac{\pi}{3} + 2\sqrt{3}
 \end{aligned}$$




$$\begin{aligned}
 14. A &= 2 \int_0^2 [2x^2 - (x^4 - 2x^2)] dx \\
 &= 2 \int_0^2 (4x^2 - x^4) dx \\
 &= 2 \left[\frac{4}{3}x^3 - \frac{1}{5}x^5 \right]_0^2 = \frac{128}{15} \approx 8.5333
 \end{aligned}$$




12. Point of intersection is given by:

$$\begin{aligned}
 x^3 - x^2 + 4x - 3 &= 0 \Rightarrow x \approx 0.783. \\
 A &\approx \int_0^{0.783} (3 - 4x + x^2 - x^3) dx \\
 &= \left[3x - 2x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 \right]_0^{0.783} \\
 &\approx 1.189
 \end{aligned}$$



$$16. y = \sqrt{x-1} \Rightarrow x = y^2 + 1$$

$$\begin{aligned}
 y &= \frac{x-1}{2} \Rightarrow x = 2y + 1 \\
 A &= \int_0^2 [(2y+1) - (y^2+1)] dy \\
 &= \int_1^5 \left[\sqrt{x-1} - \frac{x-1}{2} \right] dx \\
 &= \left[\frac{2}{3}(x-1)^{3/2} - \frac{1}{4}(x-1)^2 \right]_1^5 = \frac{4}{3}
 \end{aligned}$$




18. $A = \int_0^1 2 \, dx + \int_1^5 [2 - \sqrt{x-1}] \, dx$


$$x = y^2 + 1$$

$$A = \int_0^2 (y^2 + 1) \, dy$$

$$= \left[\frac{1}{3}y^3 + y \right]_0^2 = \frac{14}{3}$$



20. (a) $R_1(t) = 5.2834(1.2701)^t = 5.2834 e^{0.2391t}$




(b) $R_2(t) = 10 + 5.28 e^{0.2t}$

$$\text{Difference} = \int_{10}^{15} [R_1(t) - R_2(t)] \, dt \approx 171.25 \text{ billion dollars}$$


22. (a) Shell

$$V = 2\pi \int_0^2 y^3 \, dy = \left[\frac{\pi}{2}y^4 \right]_0^2 = 8\pi$$




(b) Shell

$$\begin{aligned} V &= 2\pi \int_0^2 (2-y)y^2 \, dy \\ &= 2\pi \int_0^2 (2y^2 - y^3) \, dy \\ &= 2\pi \left[\frac{2}{3}y^3 - \frac{1}{4}y^4 \right]_0^2 = \frac{8\pi}{3} \end{aligned}$$




(c) Disk

$$V = \pi \int_0^2 y^4 \, dy = \left[\frac{\pi}{5}y^5 \right]_0^2 = \frac{32\pi}{5}$$




(d) Disk

$$\begin{aligned} V &= \pi \int_0^2 [(y^2 + 1)^2 - 1^2] \, dy \\ &= \pi \int_0^2 (y^4 + 2y^2) \, dy \\ &= \pi \left[\frac{1}{5}y^5 + \frac{2}{3}y^3 \right]_0^2 = \frac{176\pi}{15} \end{aligned}$$




24. (a) Shell


$$\begin{aligned} V &= 4\pi \int_0^a (x) \frac{b}{a} \sqrt{a^2 - x^2} dx \\ &= \frac{-2\pi b}{a} \int_0^a (a^2 - x^2)^{1/2} (-2x) dx \\ &= \left[\frac{-4\pi b}{3a} (a^2 - x^2)^{3/2} \right]_0^a = \frac{4}{3}\pi a^2 b \end{aligned}$$

**(b) Disk**


$$\begin{aligned} V &= 2\pi \int_0^a \frac{b^2}{a^2} (a^2 - x^2) dx \\ &= \frac{2\pi b^2}{a^2} \left[a^2 x - \frac{1}{3} x^3 \right]_0^a \\ &= \frac{4}{3}\pi ab^2 \end{aligned}$$

**26. Disk**


$$\begin{aligned} V &= 2\pi \int_0^1 \left[\frac{1}{\sqrt{1+x^2}} \right]^2 dx \\ &= \left[2\pi \arctan x \right]_0^1 \\ &= 2\pi \left(\frac{\pi}{4} - 0 \right) \\ &= \frac{\pi^2}{2} \end{aligned}$$

**28. Disk**


$$\begin{aligned} V &= \pi \int_0^1 (e^{-x})^2 dx \\ &= \pi \int_0^1 e^{-2x} dx = \left[-\frac{\pi}{2} e^{-2x} \right]_0^1 \\ &= \left(-\frac{\pi}{2} + \frac{\pi}{2} \right) = \frac{\pi}{2} \left(1 - \frac{1}{e^2} \right) \end{aligned}$$

**30. (a) Disk**

$$\begin{aligned} V &= \pi \int_{-1}^0 x^2(x+1) dx \\ &= \pi \int_{-1}^0 (x^3 + x^2) dx \\ &= \pi \left[\frac{x^4}{4} + \frac{x^3}{3} \right]_{-1}^0 = \frac{\pi}{12} \end{aligned}$$

**(b) Shell**

$$\begin{aligned} u &= \sqrt{x+1} \\ x &= u^2 - 1 \\ dx &= 2u du \\ V &= 2\pi \int_{-1}^0 x^2 \sqrt{x+1} dx \\ &= 4\pi \int_0^1 (u^2 - 1)^2 u^2 du \\ &= 4\pi \int_0^1 (u^6 - 2u^4 + u^2) du \\ &= 4\pi \left[\frac{1}{7}u^7 - \frac{2}{5}u^5 + \frac{1}{3}u^3 \right]_0^1 = \frac{32\pi}{105} \end{aligned}$$



32. $A(x) = \frac{1}{2}bh = \frac{1}{2}(2\sqrt{a^2 - x^2})(\sqrt{3}\sqrt{a^2 - x^2})$


$$= \sqrt{3}(a^2 - x^2)$$

$$V = \sqrt{3} \int_{-a}^a (a^2 - x^2) dx = \sqrt{3} \left[a^2x - \frac{x^3}{3} \right]_{-a}^a$$

$$= \sqrt{3} \left(\frac{4a^3}{3} \right)$$

Since $(4\sqrt{3}a^3)/3 = 10$, we have $a^3 = (5\sqrt{3})/2$. Thus,

$$a = \sqrt[3]{\frac{5\sqrt{3}}{2}} \approx 1.630 \text{ meters.}$$



- 36.** Since $f(x) = \tan x$ has $f'(x) = \sec^2 x$, this integral represents the length of the graph of $\tan x$ from $x = 0$ to $x = \pi/4$. This length is a little over 1 unit. Answers (b).

34. $y = \frac{x^3}{6} + \frac{1}{2x}$

$$y' = \frac{1}{2}x^2 - \frac{1}{2x^2}$$

$$1 + (y')^2 = \left(\frac{1}{2}x^2 + \frac{1}{2x^2} \right)^2$$

$$s = \int_1^3 \left(\frac{1}{2}x^2 + \frac{1}{2x^2} \right) dx = \left[\frac{1}{6}x^3 - \frac{1}{2x} \right]_1^3 = \frac{14}{3}$$

40. $F = kx$

$$50 = k(9) \implies k = \frac{50}{9}$$

$$F = \frac{50}{9}x$$

$$\begin{aligned} W &= \int_0^9 \frac{50}{9}x dx = \left[\frac{25}{9}x^2 \right]_0^9 \\ &= 225 \text{ in} \cdot \text{lb} = 18.75 \text{ ft} \cdot \text{lb} \end{aligned}$$

38. $y = 2\sqrt{x}$

$$y' = \frac{1}{\sqrt{x}}$$

$$1 + (y')^2 = 1 + \frac{1}{x} = \frac{x+1}{x}$$

$$S = 2\pi \int_0^3 2\sqrt{x} \sqrt{\frac{x+1}{x}} dx = 4\pi \int_0^3 \sqrt{x+1} dx$$

$$= 4\pi \left[\left(\frac{2}{3} \right)(x+1)^{3/2} \right]_0^3 = \frac{56\pi}{3}$$

- 42.** We know that

$$\frac{dV}{dt} = \frac{4 \text{ gal/min} - 12 \text{ gal/min}}{7.481 \text{ gal/ft}^3} = -\frac{8}{7.481} \text{ ft}^3/\text{min}$$

$$V = \pi r^2 h = \pi \left(\frac{1}{9} \right) h$$

$$\frac{dV}{dt} = \frac{\pi}{9} \left(\frac{dh}{dt} \right)$$

$$\frac{dh}{dt} = \frac{9}{\pi} \left(\frac{dV}{dt} \right) = \frac{9}{\pi} \left(-\frac{8}{7.481} \right) \approx -3.064 \text{ ft/min.}$$

Depth of water: $-3.064t + 150$

Time to drain well: $t = \frac{150}{3.064} \approx 49 \text{ minutes}$

$(49)(12) = 588$ gallons pumped

Volume of water pumped in Exercise 41: 391.7 gallons

$$\frac{391.7}{52\pi} = \frac{588}{x\pi}$$

$$x = \frac{588(52)}{391.7} \approx 78$$

Work $\approx 78\pi \text{ ft} \cdot \text{ton}$

44. (a) Weight of section of cable: $4 \Delta x$

Distance: $200 - x$

$$W = 4 \int_0^{200} (200 - x) dx = \left[-2(200 - x)^2 \right]_0^{200} = 80,000 \text{ ft} \cdot \text{lb} = 40 \text{ ft} \cdot \text{ton}$$

- (b) Work to move 300 pounds 200 feet vertically: $200(300) = 60,000 \text{ ft} \cdot \text{lb} = 30 \text{ ft} \cdot \text{ton}$

Total work = work for drawing up the cable + work of lifting the load

$$= 40 \text{ ft} \cdot \text{ton} + 30 \text{ ft} \cdot \text{ton} = 70 \text{ ft} \cdot \text{ton}$$

46. $W = \int_a^b F(x) dx$

$$F(x) = \begin{cases} -(2/9)x + 6, & 0 \leq x \leq 9 \\ -(4/3)x + 16, & 9 \leq x \leq 12 \end{cases}$$

$$W = \int_0^9 \left(-\frac{2}{9}x + 6 \right) dx + \int_9^{12} \left(-\frac{4}{3}x + 16 \right) dx$$

$$= \left[-\frac{1}{9}x^2 + 6x \right]_0^9 + \left[-\frac{2}{3}x^2 + 16x \right]_9^{12}$$

$$= (-9 + 54) + (-96 + 192 + 54 - 144) = 51 \text{ ft} \cdot \text{lbs}$$

48. $A = \int_{-1}^3 [(2x + 3) - x^2] dx = \left[x^2 + 3x - \frac{1}{3}x^3 \right]_{-1}^3 = \frac{32}{3}$


$$\frac{1}{A} = \frac{3}{32}$$

$$\bar{x} = \frac{3}{32} \int_{-1}^3 x(2x + 3 - x^2) dx = \frac{3}{32} \int_{-1}^3 (3x + 2x^2 - x^3) dx = \frac{3}{32} \left[\frac{3}{2}x^2 + \frac{2}{3}x^3 - \frac{1}{4}x^4 \right]_{-1}^3 = 1$$

$$\bar{y} = \left(\frac{3}{32} \right) \frac{1}{2} \int_{-1}^3 [(2x + 3)^2 - x^4] dx = \frac{3}{64} \int_{-1}^3 (9 + 12x + 4x^2 - x^4) dx$$

$$= \frac{3}{64} \left[9x + 6x^2 + \frac{4}{3}x^3 - \frac{1}{5}x^5 \right]_{-1}^3 = \frac{17}{5}$$

$$(\bar{x}, \bar{y}) = \left(1, \frac{17}{5} \right)$$



50. $A = \int_0^8 \left(x^{2/3} - \frac{1}{2}x \right) dx = \left[\frac{3}{5}x^{5/3} - \frac{1}{4}x^2 \right]_0^8 = \frac{16}{5}$

$$\frac{1}{A} = \frac{5}{16}$$

$$\bar{x} = \frac{5}{16} \int_0^8 x \left(x^{2/3} - \frac{1}{2}x \right) dx$$

$$= \frac{5}{16} \left[\frac{3}{8}x^{8/3} - \frac{1}{6}x^3 \right]_0^8 = \frac{10}{3}$$

$$\bar{y} = \left(\frac{5}{16} \right) \frac{1}{2} \int_0^8 \left(x^{4/3} - \frac{1}{4}x^2 \right) dx$$

$$= \frac{1}{2} \left(\frac{5}{16} \right) \left[\frac{3}{7}x^{7/3} - \frac{1}{12}x^3 \right]_0^8 = \frac{40}{21}$$

$$(\bar{x}, \bar{y}) = \left(\frac{10}{3}, \frac{40}{21} \right)$$

